

LETTERS TO THE EDITOR

2. SERAFINI AN, WEINSTEIN MB: Case report: Radionuclide evaluation of a carotid body tumor. *J Nucl Med* 13: 640-643, 1972

3. DESAI MG, PATEL CC: Heredo-familial carotid body

tumors. *Clin Radiol* 12: 214-218, 1961

4. KORN D, BENSCH K, LIEBOW AA, et al: Multiple minute pulmonary tumors resembling chemodectomas. *Am J Pathol* 37: 641-672, 1960

REPLY

Drs. Moinuddin and Rockett call attention to the interesting parallel between our static images of a chemodectoma obtained with ^{99m}Tc -macroaggregated albumin and previously reported dynamic images of chemodectomas obtained with sodium pertechnetate. We do not think the pulmonary lesions in our case were chemodectomas since our reported findings show that the pulmonary lesions passed the albumin macroaggregates while the cervical lesion retained them. Furthermore, the pulmonary chemodectomas described by Korn et al, cited by Moinuddin and Rockett, were only large enough to be visible without the microscope in one case out of nineteen. In that one case, no lesions larger than 3 mm were described.

The lesions in our case, on the other hand, ranged up to several centimeters in size. Their plain chest film appearance was very suggestive of arteriovenous malformations. The pulmonary angiogram showed typical arteriovenous malformations with large feeding arteries and draining veins and no tumor vascularity. Hence, it is extremely unlikely that these lesions represent anything but arteriovenous malformations.

C. D. RUSSELL

H. P. JANDER

E. V. DUBOVSKY

University of Alabama Medical Center
Birmingham, Alabama

GALLIUM-67 UPTAKE IN THE REGENERATING RAT LIVER

Recently, Hill and Wagner (1) have reported that the uptake of ^{67}Ga in regenerating liver is similar to that in normal liver; they suggest that " ^{67}Ga uptake is not related to hepatic cell proliferation associated with regeneration." While we do not disagree with this conclusion, we have observed considerable variation over the 72 hr after partial hepatectomy in ^{67}Ga concentration in the regenerating liver. Studies were taken 2 hr after intravenous injection of ^{67}Ga -citrate (2). Gallium-67 uptake was found to be maximal (about four times control levels) at 42 hr after operation and minimal (approximately 1.4 times control levels) during the period of stimulated DNA synthesis. Hill and Wagner confined their studies to this latter period.

In vitro studies with synchronized cultures of HeLa cells have also shown that ^{67}Ga uptake reaches a nadir at the time of most rapid DNA synthesis; maximum uptake of the nuclide was observed in the G_2 phase of the cell cycle.

Thus, while ^{67}Ga uptake does not appear to be directly related to cell proliferation per se, there do appear to be significant variations in the nuclide uptake at different stages of the cell cycle. In regenerating rat liver, there is a good correlation between variations in ^{67}Ga uptake and in lysosomal enzyme activity during the early regeneration period.

PETER A. G. HAMMERSLEY

MAUREEN A. ZIVANOVIC

Institute of Cancer Research
The Royal Marsden Hospital
Sutton, Surrey, England

REFERENCES

1. HILL JH, WAGNER HN: ^{67}Ga -uptake in the regenerating rat liver. *J Nucl Med* 15: 818-820, 1974

2. HAMMERSLEY PAG, CAUCHI MN, TAYLOR DM: Gallium uptake in the regenerating rat liver, and its relationship to lysosomal enzyme activity. *Cancer Res* 35: 1154-1158, 1975

TECHNETIUM LABELING OF STREPTOKINASE AT LOW pH

Having read the paper by Persson and Kampi (1) with interest, we would like to make a number of observations from our experience. Like Persson and Kampi, we found improved labeling of ^{99m}Tc -streptokinase at low pH values (pH 2-3). Labeling at high pH (pH 12) using a modification of the method of Dugan et al (2) gave labeling yields of more than 10%. Since the standard deviation of our analytic result was high with the G-25 gel-chromatography

quality-control method, we preferred thin-layer chromatography (protein localization with ninhydrin combined with radiochromatogram scanning). The enzyme activity (activating plasminogen to plasmin) of ^{99m}Tc -streptokinase was found to be decreased at high pH.

In experiments with rabbits, ^{99m}Tc -streptokinase gave good results. However, the problem of antibody