

Rev Clin Radiol Nucl Med 5: 523-558, 1974

3. ANDERSON JT, DEHNER LP: Osteolytic form of Paget's disease. *J Bone Joint Surg* 58A: 994-1000, 1976
4. LENTLE BC, RUSSELL AS, HERSLIP PG, et al: The scintigraphic findings in Paget's disease of bone. *Clin Radiol* 27: 129-135, 1976
5. FRANCIS MD: The inhibition of calcium hydroxyapatite crystal growth by polyphosphonates and polyphosphates. *Calcif Tissue Res* 3: 151-162, 1969

Splenic Uptake of Tc-99m Sulphur Colloid in Malignant Melanoma

Goldman (1) reported the observation that ten of 22 patients with malignant melanoma, in whom no hepatic abnormality could be detected by scan or liver chemistry, showed augmented splenic radioactivity in Tc-99m sulphur colloid scans. In the same report, none of 60 patients with carcinoma at various sites demonstrated such an increase in splenic uptake. Anterior and left anterior oblique scans of the spleen were carried out using a rectilinear scanner, and counts per minute were recorded over the area of highest activity. Rates of particle extraction in normal patients were demonstrated graphically by additional printouts stored on magnetic tapes during sequential gamma-camera liver imaging over a 15-min period. Criteria for increased splenic uptake included splenic density exceeding that of the liver on the anterior projection, and more counts per minute recorded over splenic "hot spots" on the left anterior oblique scan of the liver, compared with the anterior scan.

Shortly thereafter Klingensmith (2) described a patient in whom increased splenic uptake (using criteria similar to those of the Goldman study) was noted. In this patient the abnormal increase in spleen density disappeared following resection of a primary melanoma that was confined to the globe of the eye.

Both investigators theorized that malignant melanoma may produce a factor that increases uptake of Tc-99m sulphur colloid in the reticuloendothelial cells of the spleen.

We have recently reviewed our experience with 34 scans in 24 patients who were put on an adjuvant immunotherapy program for malignant melanoma, with the presumption that no distant or visceral disease was present. In all of these patients at least one scan was done before the onset of treatment.

In only two patients was increased splenic uptake noted. One was a patient in whom occult hepatic disease, which was not previously appreciated, was seen on liver scan. In the second such patient, demonstration of increased splenic uptake was followed at 5 mo by a scan with decrease in overall density of uptake in the spleen, but with the appearance of a discrete filling defect presumably representing a metastatic lesion. Four of these 24 patients had uptake considered to be at the upper limit of normal, and one patient demonstrated "small normal" category of splenic uptake.

An additional patient of interest was seen separately from this study. In this patient, also on immunotherapy, diffuse enlargement and increase in density of spleen was followed by splenectomy, which demonstrated what was initially thought to be "Hairy Cell" leukemia by the pathologist who saw the slides. It was later shown that this diffuse splenic enlargement was in fact due to a miliary distribution of splenic metastases from melanoma.

In summary, in a brief review of 24 patients in whom only primary and/or regional melanoma was thought to be present, no patients were found in whom splenic enlargement or increased density of uptake, demonstrated by spleen scan, could not be explained by probable splenic metastases. This is clearly contrary to the experience of Sober (3) and others who have recently reported identification of 50 patients with increased splenic uptake on scan in a series of 148 patients with early melanoma without known visceral disease. Only one of 165 patients with other malignancies, in their experience, demonstrated such an increase in splenic uptake.

Our findings suggest that excess splenic uptake observed in melanoma is either a nonspecific phenomenon related to previously unappreciated diffuse hepatic disorder (e.g., cirrhosis, hepatitis, or metastases) or nonspecific stimulation of the RE system (e.g., by septicemia). An additional explanation is that it may relate to the technical problem of estimating absolute counting rates in three-dimensional organs that are looked at in an essentially two-dimensional pattern. If true increase in splenic uptake of the sulphur colloid does occur, it may be a relatively uncommon event, probably not useful for diagnostic or prognostic purposes.

LARRY NATHANSON

PAUL KAHN

Tufts-New England Medical Center
Boston, Massachusetts

REFERENCES

1. GOLDMAN AB, BRAUNSTEIN P, SONG C: Augmented splenic uptake of ^{99m}Tc-sulphur colloid in patients with malignant melanoma. *Radiology* 112: 631-634, 1974
2. KLINGENSMITH WC: Resolution of increased splenic uptake and uptake of ^{99m}Tc-Sulphur Colloid following removal of a malignant melanoma. *J Nucl Med* 15: 1203-1204, 1974
3. SOBER A, et al: Minutes of the Melanoma Discussion Group, January 18, 1977. Bethesda, Maryland, NIH, p 7

Skeletal Uptake of Tc-99m HEDP in Primary Hyperparathyroidism

Using Tc-99m labeled phosphate compounds, bone scanning and simple quantification have been performed in patients with hyperparathyroidism, but contradictory results have been obtained (1-3).

Sy (1) described four patients with primary hyperparathyroidism whose scans with Tc-99m polyphosphate showed relatively increased skeletal uptake of radiopharmaceutical in calvarium, mandible and long-bone epiphyses, but he did not quantify his results. However, Weigmann et al. (2), using Tc-99m pyrophosphate, found that most of their 20 patients with primary hyperparathyroidism had apparently normal bone scans, and also their ratios of bone to soft tissue were not significantly different from a control group. Krishnamurthy et al. (3), also using Tc-99m pyrophosphate, found that seven of their 12 patients with primary hyperparathyroidism had bone-scan abnormalities that were located mainly at the extremities. They also measured urinary excretion of Tc-99m pyrophosphate in nine of those patients over a 4-hr period and found this to be higher than in a control group. This implies that (unless soft-tissue uptake is altered) total skeletal uptake of the radiopharmaceutical is reduced in such cases.