

- J, Huang SC, Schelbert HR. Quantification of absolute myocardial perfusion at rest and during exercise with positron emission tomography after human cardiac transplantation. *J Am Coll Cardiol* 1991;18:512-517.
49. Rechavia E, Araujo LI, De Silva R, et al. Dipyridamole vasodilator response after human orthotopic heart transplantation: quantification by oxygen-15-labeled water and positron emission tomography. *J Am Coll Cardiol* 1992;19:100-106.
50. Hoff SJ, Stewart JR, Frist WH, et al. Noninvasive detection of heart transplant rejection with positron emission scintigraphy. *Ann Thorac Surg* 1992;53:572-577.
51. Schwaiger M, Kalff V, Rosenspire K, et al. Noninvasive evaluation of sympathetic nervous system in human heart by positron emission tomography. *Circulation* 1990;82:457-464.
52. Schwaiger M, Hutchins GD, Kalff V, et al. Evidence for regional catecholamine uptake and storage sites in the transplanted human heart by positron emission tomography. *J Clin Invest* 1991;87:1681-1690.

ERRATUM

Due to a production error, Figure 1 in the article, "Reproducibility of Repeated Measures of Carbon-11-Raclopride Binding in the Human Brain," by Volkow et al. in the April issue of the *Journal* (p. 611) was printed incorrectly. The corrected figure and legend are shown below.




FIGURE 1. Location, size and shape of ROIs used to quantitate ^{11}C -raclopride in basal ganglia (upper images) and cerebellum (lower image).