
A Kit Formulated Under Good Manufacturing Practices for Labeling Human Epidermal Growth Factor with ^{111}In for Radiotherapeutic Applications

Raymond M. Reilly, PhD¹⁻³; Deborah A. Scollard, BAppSci¹; Judy Wang, MSc¹; Hridya Mondal, BSc¹; Paul Chen, PhD¹; Lee A. Henderson, PhD⁴; Barry M. Bowen, MSc²; and Katherine A. Vallis, MD, PhD^{5,6}

¹Division of Nuclear Medicine, Toronto General Hospital, University Health Network, Toronto, Ontario, Canada; ²Department of Pharmaceutical Sciences, University of Toronto, Toronto, Ontario, Canada; ³Department of Medical Imaging, University of Toronto, Toronto, Ontario, Canada; ⁴Viral Therapeutics Inc., Ithaca, New York; ⁵Department of Radiation Oncology, Princess Margaret Hospital, University Health Network, Toronto, Ontario, Canada; and ⁶Department of Medical Biophysics, University of Toronto, Toronto, Canada

Our goal was to design and manufacture a kit under good manufacturing practices (GMP) for the preparation of ^{111}In -DTPA-hEGF Injection, a novel targeted radiotherapeutic agent for advanced epidermal growth factor receptor (EGFR)-positive breast cancer. **Methods:** Human EGF (hEGF) was derivatized with diethylenetriaminepentaacetic acid (DTPA) and then purified by size-exclusion chromatography and ultrafiltration. Kits were prepared by dispensing 0.25 mg (1 mL) of DTPA-hEGF in 1 mol/L sodium acetate buffer [pH 6.0] into single-dose glass vials. Raw materials were pharmacopoeial or reagent grade according to the American Chemical Society and were tested for identity and purity. Kits were tested for protein concentration, purity and homogeneity (sodium dodecyl sulfate polyacrylamide gel electrophoresis and size-exclusion high-performance liquid chromatography), pH, clarity and color, volume, DTPA substitution, labeling efficiency, receptor binding to MDA-MB-468 human breast cancer cells, and sterility and apyrogenicity. ^{111}In -DTPA-hEGF Injection was tested for pH, radionuclidic and radiochemical purity, clarity and color, and sterility and apyrogenicity. **Results:** Four lots of kits and 8 lots of ^{111}In -DTPA-hEGF Injection passed all quality specifications. The labeling efficiency was 94%–99% with 115–773 MBq ^{111}In chloride added to a single kit. ^{111}In -DTPA-hEGF exhibited preserved receptor binding against MDA-MB-468 cells (affinity constant [K_a], $0.9\text{--}1.1 \times 10^7$ L/mol; maximum number of binding sites per cell [B_{max}], $1.1\text{--}2.2 \times 10^6$ sites per cell). In addition, labeling of aliquots of the kit suggested that a single vial could be labeled with up to 3,083 MBq ^{111}In while maintaining a radiochemical purity of >90%. Kits were stable for >90 d and ^{111}In -DTPA-hEGF Injection was stable for >24 h stored at 4°C. **Conclusion:** The kit formulation is suitable for preparing ^{111}In -DTPA-hEGF Injection for a phase I clinical trial in patients with advanced EGFR-positive breast cancer. Establishment of the GMP pro-

cesses for ^{111}In -DTPA-hEGF Injection provides a useful example of manufacturing biotechnology-based investigational radiopharmaceuticals in an academic environment for early phase I clinical trials.

Key Words: Auger electrons; ^{111}In ; epidermal growth factor; kit; good manufacturing practices

J Nucl Med 2004; 45:701–708

Radiopharmaceuticals labeled with low-energy Auger electron-emitting radionuclides (e.g., ^{111}In or ^{125}I) are receiving considerable interest as targeted radiotherapeutic agents for cancer (1). Theoretically, the micrometer range of the electrons should restrict their radiotoxicity mainly toward cells that internalize the radiopharmaceuticals into the cytoplasm and especially in cases in which they are translocated to the cell nucleus (2,3). Our laboratory has discovered a novel targeted Auger electron-emitting radiopharmaceutical, ^{111}In -labeled human epidermal growth factor (^{111}In -DTPA-hEGF) (where DTPA is diethylenetriaminepentaacetic acid) (4), which exploits the overexpression of epidermal growth factor receptors (EGFRs) present on almost all estrogen receptor-negative, hormone-resistant, and poor-prognosis breast cancers (5).

^{111}In -DTPA-hEGF was rapidly internalized into the cytoplasm and translocated to the nucleus of EGFR-positive human breast cancer cells, where the Auger electron emissions were highly damaging to DNA, causing cell death (4). The radiopharmaceutical was highly cytotoxic to MDA-MB-468 human breast cancer cells overexpressing EGFR ($1\text{--}2 \times 10^6$ receptors per cell) with >95% cell killing achieved at <111–148 mBq per cell (4). Furthermore, the radiopharmaceutical was 85–300 times more potent at inhibiting the growth of MDA-MB-468 cells in vitro than the

Received Sep. 3, 2003; revision accepted Dec. 2, 2003.

For correspondence or reprints contact: Raymond M. Reilly, PhD, Leslie Dan Faculty of Pharmacy, University of Toronto, 19 Russell St., Toronto, ON, Canada, M5S 2S2.

E-mail: raymond.reilly@utoronto.ca.

chemotherapeutic agents methotrexate, doxorubicin, and paclitaxel and several orders of magnitude more effective than 5-fluorouracil (6). Administration of 5 weekly doses (18.5 MBq; 3 μ g) of ^{111}In -DTPA-hEGF to athymic mice caused growth arrest of established subcutaneous MDA-MB-468 xenografts with minimal normal tissue toxicity (modest decrease in leukocyte and platelet counts) (7). Early treatment of mice bearing smaller "nonestablished" MDA-MB-468 xenografts with ^{111}In -DTPA-hEGF achieved tumor regression.

To translate ^{111}In -DTPA-hEGF from preclinical investigation to a phase I clinical trial, it is necessary to create a pharmaceutical quality formulation manufactured under current Good Manufacturing Practices (GMP) and obtain regulatory approval from Health Canada in the form of a Clinical Trial Application (CTA). GMP are comprehensive quality processes that ensure the suitability of the product for its intended use in humans (8). Meeting GMP requirements is one of the major challenges facing radiopharmaceutical scientists who conduct translational research and work at a university or hospital setting with limited resources. In this study, we describe our approach to manufacturing a kit for the preparation of ^{111}In -DTPA-hEGF Injection under GMP in the clinical radiopharmaceutical research laboratory at the University Health Network, a University of Toronto-affiliated hospital. We propose that the strategy for establishing GMP for ^{111}In -DTPA-hEGF Injection provides a useful example of manufacturing biotechnology-based investigational radiopharmaceuticals in an academic environment for early phase I clinical trials.

MATERIALS AND METHODS

Raw Materials

hEGF (>98%) was obtained as hEGF₁₋₅₃ from Upstate Biotechnology Inc. or as hEGF₁₋₅₁ from Viral Therapeutics Inc. DTPA dianhydride (>98%) and chloroform (reagent grade, >99.9% meeting specifications of the American Chemical Society) (9) were purchased from Sigma-Aldrich Canada Ltd. Sodium acetate dihydrate USP and sodium bicarbonate USP were obtained from EM Science. Sterile Water for Injection USP and Sodium Chloride Injection USP were obtained from BaxterTravenol Inc. Nitrogen NF was obtained from Praxair Canada, Inc. All other chemicals and reagents were purchased in analytic grade with a minimum purity of >95%. Sterile, apyrogenic type 1 glass multidose vials (10 mL) with a gray butyl rubber septum and aluminum seal were obtained from Hollister-Stier Laboratories Inc. ^{111}In chloride (>3.7 GBq/mL; <0.1% $^{114\text{m}}\text{In}$ and ^{65}Zn) was purchased in radiochemical quality from MDS Nordion Inc. or PerkinElmer Life Sciences Inc.

Identity Testing and Purity Assessment of Raw Materials

Certificates of analysis were obtained from the vendor for each lot of raw materials. Identity testing of sodium bicarbonate USP and sodium acetate USP was performed by pharmacopoeial methods (10,11). The purity of nonpharmacopoeial materials was confirmed by in-house analytic techniques. Proton (^1H) NMR (500 MHz) spectroscopy was used to confirm the identity of chloroform

(neat) and DTPA dianhydride (dissolved in D_2O). The purity of DTPA dianhydride was measured by adapting the assay for Edetic Acid NF (12) using a 10.0 mg/mL solution of the raw material to titrate a known amount of chelometric standard calcium carbonate (100.00%; Fisher Scientific Ltd.). The identity and radionuclidic purity of ^{111}In chloride was confirmed by γ -spectroscopy on a Capintec model 2000 multichannel analyzer (Capintec, Inc.) checked using radionuclide disk reference sources (^{133}Ba , ^{22}Na , ^{137}Cs , and ^{60}Co) and with a certified primary reference standard for ^{111}In (National Institute of Standards and Technology).

Characterization and Purity Evaluation of hEGF

Amino acid analysis, ultraviolet (UV) spectroscopy, and sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE)/Western blot were used to characterize hEGF, and size-exclusion high-performance liquid chromatography (HPLC) was used to measure its purity and homogeneity. UV spectroscopy was performed for hEGF₁₋₅₃ (0.25 mg/mL in 50 mmol/L sodium bicarbonate buffer, pH 7.5). SDS-PAGE was conducted on a 4%–20% Tris HCl gradient minigel (Bio-Rad Laboratories, Inc.) stained with Coomassie R-250 brilliant blue. Western blot was performed by transferring electrophoresed proteins onto a nitrocellulose membrane (TransBlot; Bio-Rad Laboratories) and probing with a rabbit polyclonal anti-hEGF antibody (provided by Dr. Jean Gariépy, Ontario Cancer Institute). Reactive bands were detected with a goat antirabbit IgG-horseradish peroxidase conjugate (Sigma-Aldrich Canada Ltd.) and diaminobenzidine/0.03% H_2O_2 . Size-exclusion HPLC was performed on a BioSep SEC-S2000 column (Phenomenex Inc.) eluted with 100 mmol/L NaH_2PO_4 buffer (pH 7.0) at a flow rate of 1.0 mL/min using a Beckman System Gold model 125 HPLC interfaced with a model 166 UV detector (Beckman Coulter) set at 280 nm.

Pharmaceutical Buffers

Sterile, nonpyrogenic 50 mmol/L sodium bicarbonate (pH 7.5) in Sodium Chloride Injection USP and 1 mol/L sodium acetate buffer (pH 6.0) (in Sterile Water for Injection USP) buffers were prepared from pharmacopoeial-quality raw materials. Trace metals were stripped from the buffers by passage through a cation-exchange column consisting of a 60-mL sterile syringe plugged with glass wool and filled with 30 mL of Chelex-100 resin (Bio-Rad Laboratories) prehydrated overnight in Sterile Water for Injection USP. After removal of trace metals, the pH was readjusted to the desired value using sterile 1N HCl and the buffers were sterilized by filtration through a 0.22- μm pore size Millex-GV filter (Millipore Corp.). Quality control testing included USP Sterility and Pyrogen Tests as well as an assay for the concentration of sodium acetate or sodium bicarbonate by USP methods (10,11). The assay for sodium acetate consisted of titration of the buffer with standardized 0.1N perchloric acid (Fisher Scientific Ltd.). The assay for sodium bicarbonate consisted of titration with standardized 0.1N sulfuric acid (Fisher Scientific Ltd.). The stability of the buffers stored at 4°C was determined by reassaying the concentration of sodium acetate or sodium bicarbonate up to 11 mo after preparation.

Radiopharmaceutical Kits

A kit for the preparation of ^{111}In -DTPA-hEGF Injection was constructed by derivatizing hEGF with a 20-fold molar excess of DTPA dianhydride. Briefly, DTPA dianhydride (50 mg) was suspended in 5.0 mL of chloroform in a sterilized 10-mL glass scintillation vial, and a 600- μL aliquot (17 μmol) was dispensed

into a sterilized 10-mL glass Reacti-Vial (Pierce Biotechnology, Inc.). Additional chloroform was added to a final volume of 1.0 mL; then the chloroform was evaporated to dryness using a gentle stream of nitrogen NF. Approximately 1.0 mL (5 mg; 0.83 μ mol) of hEGF in 50 mmol/L sodium bicarbonate buffer (pH 7.5) was added and the vial was vortexed for 1 min. The reaction mixture was allowed to incubate at room temperature for a further 30 min. A 10- μ L aliquot of the reaction mixture was removed for measurement of DTPA conjugation efficiency while the remainder was transferred to the top of a 1 \times 20 cm P-2 size-exclusion chromatography column (exclusion limit, 1.8 kDa; Bio-Rad). The column was eluted with 20 \times 0.5-mL aliquots of 50 mmol/L sodium bicarbonate buffer (pH 7.5), and the fractions were collected in sterile, polystyrene tubes (VWR International). The absorbance of each fraction was measured at 280 nm. The fractions containing the partially purified DTPA-hEGF (usually fractions 5–12) were combined. The pooled fractions were transferred in 2 equal portions to Centricon YM-3 ultrafiltration devices (molecular weight [M_r] cutoff = 3 kDa; Millipore Corp.), and the solution in each device was diluted to 2.0 mL with 1 mol/L sodium acetate buffer (pH 6.0). The Centricon YM-3 devices were centrifuged at 4,500 rpm (2,875g) for 45 min in a fixed-angle centrifuge (model Centra-4B; IEC). The solutions were rediluted to 2.0 mL with 1 mol/L sodium acetate buffer (pH 6.0) and the devices were recentrifuged. A total of 8 dilution and ultrafiltration steps were performed. Finally, the pure DTPA-hEGF solutions were recovered in 0.5-mL volume and combined. The concentration of DTPA-hEGF was assayed spectrophotometrically at 280 nm by reference to a calibration curve created using hEGF_{1–53} standards (0–0.5 mg/mL). DTPA-hEGF was diluted to a final concentration of 0.25 mg/mL with 1 mol/L sodium acetate buffer (pH 6.0) and sterilized by filtration through a 0.22- μ m Millex-GV filter. Unit-dose radiopharmaceutical kits were prepared by aseptically dispensing 1.0-mL (0.25 mg) aliquots into sterile, apyrogenic 10-mL glass unit-dose vials using a 1-mL sterile syringe and needle in a laminar flow hood.

Quality Control Testing of Kits

The pharmaceutical quality of the kits was evaluated by determining the protein concentration, protein homogeneity and polymerization, pH, clarity and color, volume contained in each vial, DTPA substitution level, labeling efficiency with ¹¹¹In, receptor-binding properties, and sterility and apyrogenicity. The concentration of hEGF was measured spectrophotometrically at 280 nm. Protein homogeneity and polymerization were evaluated by SDS-PAGE and size-exclusion HPLC. The pH was measured using narrow-range pH paper (range, 4.5–7.5 in 0.5-unit increments; Fisher Scientific Ltd.). Clarity and color were evaluated by inspection against a light or dark background under bright light. The volume of solution contained in each vial was determined by weighing the vials before and after filling, assuming a density of 1 g/mL at 20°C. DTPA conjugation efficiency was determined by trace labeling a 10- μ L aliquot (50 μ g) of the unpurified reaction mixture with 1 MBq ¹¹¹In and determining the proportion of ¹¹¹In-DTPA-hEGF and ¹¹¹In-DTPA by instant thin-layer silica gel chromatography (ITLC-SG; Pall Corporation) developed in 100 mmol/L sodium citrate (pH 5.0). R_f values for ¹¹¹In-DTPA-hEGF and ¹¹¹In-DTPA in this system were 0.0 and 1.0, respectively. The DTPA substitution level was calculated by multiplying the conjugation efficiency by the molar ratio of DTPA dianhydride to hEGF used in the reaction (i.e., 20:1).

The labeling efficiency of the kits was determined by adding 185 MBq ¹¹¹In chloride to a single vial, incubating for 30 min, and

determining the percentage of ¹¹¹In-DTPA-hEGF by ITLC-SG. The labeling efficiency of the kits using ¹¹¹In chloride from 2 different suppliers (MDS Nordion Inc. and PerkinElmer Life Sciences Inc.) was compared. The maximum amount of radioactivity that could be added to the kits while maintaining a radiochemical purity of >90% was studied by labeling 25 μ L of kit solution containing 6 μ g DTPA-hEGF with increasing amounts of ¹¹¹In chloride (1.1–74 MBq) corresponding to the addition of 46–3,083 MBq to a single vial. The stability of the kits stored at 4°C was evaluated by retesting against all specifications (except sterility and apyrogenicity) at up to 90 d after manufacture.

Measurement of Receptor-Binding Properties

The equivalence of hEGF_{1–51} and hEGF_{1–53} raw materials was evaluated by comparing their ability to displace the binding of ¹²³I-hEGF_{1–53} to MDA-MB-468 human breast cancer cells (1–2 \times 10⁶ EGFRs per cell; American Type Culture Collection). ¹²³I-hEGF was prepared as previously described (13). Briefly, ¹²³I-hEGF_{1–53} (3 ng; 3.7 mBq) was incubated for 30 min at 37°C with 1 \times 10⁶ MDA-MB-468 cells in the presence of increasing concentrations (1 nmol/L to 10 μ mol/L) of hEGF_{1–51} or hEGF_{1–53} in 150 mmol/L sodium chloride. The tubes were centrifuged and the cell pellet was separated and measured in a γ -counter. The receptor-binding curve was obtained by plotting the radioactivity bound to the cells versus the concentration of competitor (hEGF_{1–51} or hEGF_{1–53}). The dissociation constant (K_d) values were estimated by fitting the curve to a 1-site competition receptor-binding model using GraphPad Prism 3.0 software (GraphPad Software, Inc.). The receptor-binding properties of ¹¹¹In-DTPA-hEGF Injection were evaluated in a direct receptor-binding assay using MDA-MB-468 cells as previously reported (13). The affinity constant (K_d) and maximum number of binding sites per cell (B_{max}) were estimated by fitting the curve to a 1-site direct receptor-binding model using GraphPad Prism 3.0 software.

Final Radiopharmaceutical

¹¹¹In-DTPA-hEGF Injection was prepared by aseptically removing the cap from a single unit-dose vial of the kit in a laminar flow hood and adding 115–960 MBq (5–20 μ L) ¹¹¹In chloride directly into the vial using an Eppendorf micropipette and sterile pipette tip. After an incubation period of at least 30 min, the radiopharmaceutical was diluted to 3.0 mL with Sodium Chloride Injection USP. The radiopharmaceutical was drawn up in a lead-shielded syringe and sterilized by filtration through a 0.22- μ m Millex-GV filter into a 10-mL sterile, nonpyrogenic glass vial. Quality control of ¹¹¹In-DTPA-hEGF Injection included measurement of total radioactivity, pH, radiochemical purity, clarity and color, and sterility and apyrogenicity. Total radioactivity was measured in a radioisotope calibrator (Capintec model CRC-12). Radiochemical purity, pH, clarity, and color were determined as described previously. Radionuclidic purity was determined on the ¹¹¹In chloride raw material. Sterility and apyrogenicity were assessed retrospectively by USP Sterility and Pyrogen Tests after allowing 30 d for radionuclide decay. The stability of ¹¹¹In-DTPA-hEGF stored at 4°C was evaluated by measuring the radiochemical purity up to 24 h after preparation.

RESULTS

Raw Materials

All raw materials passed tests for identity and met specifications for purity (>95%). The amino acid analyses of

hEGF₁₋₅₃ and hEGF₁₋₅₁ were consistent with their known composition. The C-terminal residues Leu-52 and Arg-53 in hEGF₁₋₅₃ are not present in hEGF₁₋₅₁ nor are they required for receptor binding (14,15). The identity of hEGF was further confirmed by the absence of threonine or phenylalanine in the peptide (16). The UV spectrum of hEGF₁₋₅₃ (not shown) exhibited λ_{\max} values of 220 nm ($\epsilon = 92,700$ mol/L⁻¹) and 280 nm ($\epsilon = 18,500$ mol/L⁻¹). SDS-PAGE analysis of hEGF₁₋₅₃ or hEGF₁₋₅₁ showed 1 major band corresponding to a protein with the expected M_r of 6 kDa (Fig. 1A) and a minor closely migrating band corresponding to a protein with slightly lower M_r . The major band was positive on Western blot when probed with a rabbit polyclonal anti-hEGF antibody (Fig. 1B). Size-exclusion HPLC of hEGF₁₋₅₃ or hEGF₁₋₅₁ (not shown) demonstrated 1 major peak with a retention time (t_R) of 11.5 min. There were no major peaks in the HPLC analysis of hEGF associated with impurities indicating a purity of >95%. There was no difference in the ability of hEGF₁₋₅₃ or hEGF₁₋₅₁ to compete with ¹²⁵I-hEGF₁₋₅₃ for binding to MDA-MB-468 breast cancer cells (Fig. 2). Proton (¹H) NMR (500 MHz) spectra of DTPA dianhydride and chloroform (not shown) were consistent with their chemical structures. The purity of DTPA dianhydride (102.9%) was within specifications (95%–105%). There were no detectable ^{114m}In or ⁶⁵Zn radionuclidic impurities in ¹¹¹In chloride. The expiry of all raw materials (except ¹¹¹In chloride) was set arbitrarily at 2 y from receipt.

Pharmaceutical Buffers

Four lots of 50 mmol/L sodium bicarbonate buffer (pH 7.5) and 1 mol/L sodium acetate buffer (pH 6.0) were prepared. Each lot met specifications for sodium bicarbonate or sodium acetate, pH, clarity, and color. The buffers were sterile and pyrogen free and were stable stored at 4°C.




FIGURE 1. (A) SDS-PAGE analysis of hEGF raw materials on 4%–20% Tris HCl gradient minigel stained with Coomassie R-250 brilliant blue. MW = broad-range molecular weight markers; UB = hEGF₁₋₅₃ (Upstate Biotechnology Inc.; 2 μ g); VTI = hEGF₁₋₅₁ (Viral Therapeutics Inc.; 2 μ g). (B) Corresponding Western blot using polyclonal rabbit anti-hEGF antibody.




FIGURE 2. Competition receptor-binding assay curve for displacement of binding of ¹²⁵I-hEGF₁₋₅₃ to MDA-MB-468 human breast cancer cells by hEGF₁₋₅₃ (○) or hEGF₁₋₅₁ (●) raw materials. Each point represents a single determination. K_d values for hEGF₁₋₅₃ and hEGF₁₋₅₁ were 1.4×10^{-7} and 2.5×10^{-7} mol/L, respectively.

The concentration of sodium bicarbonate in 50 mmol/L sodium bicarbonate buffer (pH 7.5) remained within $\pm 5\%$ of the initial assay value up to 9 mo after manufacturing. The concentration of sodium acetate in 1 mol/L sodium acetate buffer (pH 6.0) similarly remained within $\pm 5\%$ of the initial assay value up to 11 mo after manufacturing. Based on the stability data, the expiry of the pharmaceutical buffers was set at 9 mo.

Radiopharmaceutical Kits

Three lots of kits for the preparation of ¹¹¹In-DTPA-hEGF Injection were prepared with hEGF₁₋₅₃ raw material (2F004, 2G004, and 2I002) and 1 lot (3B003) was prepared using hEGF₁₋₅₁ (Table 1). Each lot of kits met specifications for protein concentration, pH, clarity and color, DTPA substitution level, purity and homogeneity, labeling efficiency with ¹¹¹In, receptor binding, and sterility and apyrogenicity. SDS-PAGE analysis of DTPA-hEGF (not shown) revealed 1 major band corresponding to a protein with a M_r of 6 kDa and a second minor band corresponding to a protein with a M_r of 12 kDa, representing monomeric and dimeric DTPA-hEGF, respectively. Dimeric DTPA-hEGF is due to protein crosslinking through the DTPA moiety caused by the bifunctional nature of DTPA dianhydride. Size-exclusion HPLC (not shown) similarly demonstrated a major peak with a t_R of 11.5 min representing monomeric DTPA-hEGF and a second minor peak (<5%) with a t_R of 10.5 min representing dimeric DTPA-hEGF. The labeling of the kits with ¹¹¹In was rapid, reproducible, and almost quantitative (94%–99%; Table 1). One lot of kits (2G004) labeled with ¹¹¹In chloride (185 MBq) from 2 different

TABLE 1
Quality Control Testing of Kits for Preparation of ^{111}In -DTPA-hEGF Injection

Test	Specification	Kit lot			
		2F004	2G004	2I002	3B003
Protein concentration (mg/mL)	0.22–0.27	0.22	0.25	0.23	0.23
SDS-PAGE	1 major band at 6 kDa; 1 minor band at 12 kDa	Passed	Passed	Passed	Passed
Size-exclusion HPLC	1 major peak t_R 11.5 min; 1 minor peak t_R 10.5 min	Passed	Passed	Passed	Passed
pH	5.5–6.5	6.5	6.5	6.5	6.0
Clarity and color	Clear and colorless	Passed	Passed	Passed	Passed
Volume (mL)	0.9–1.1	0.9–1.1	0.9–1.1	0.9–1.1	0.9–1.1
DTPA substitution	0.5–1.0 DTPA/hEGF	0.53	0.50	0.52	0.76
Labeling efficiency (%)	>90	94.0	97.5	98.8	98.0
Receptor binding	$K_a = 0.5\text{--}2 \times 10^7$ L/mol; $B_{\max} = 0.5\text{--}4 \times 10^6$ sites/cell	1.1×10^7 2.0×10^6	1.1×10^7 2.2×10^6	0.9×10^7 1.1×10^6	2.2×10^7 1.1×10^6
Sterility	USP XXV	Passed	Passed	Passed	Passed
Apyrogenicity	USP XXV	Passed	Passed	Passed	Passed

suppliers (MDS Nordion Inc. and PerkinElmer Life Sciences Inc.) exhibited a labeling efficiency of 97.0% and 96.3%, respectively. The labeling efficiency for aliquots of the kit solution (25 μL ; 6 μg DTPA-hEGF) incubated with 74 MBq ^{111}In was $91.2\% \pm 0.3\%$. These results suggested that a single vial (1 mL; 250 μg DTPA-hEGF) could be labeled with up to 3,083 MBq ^{111}In and remain within specifications for radiochemical purity (>90%). ^{111}In -DTPA-hEGF Injection demonstrated specific and saturable binding to MDA-MB-468 breast cancer cells (Fig. 3). The mean K_a for ^{111}In -DTPA-hEGF was $1.3 \pm 0.6 \times 10^7$ L/mol and the B_{\max} was $1.6 \pm 0.6 \times 10^6$ sites per cell. All kits retested for quality at 90 d continued to meet specifications. There was no significant decrease in labeling efficiency at

90 d compared with initial testing values ($96.7\% \pm 1.3\%$ vs. $97.1\% \pm 2.1\%$, respectively), and there was no change in the receptor-binding properties (mean K_a , $1.7 \pm 0.6 \times 10^7$ L/mol; B_{\max} , $2.3 \pm 0.2 \times 10^6$ sites per cell). The expiry of the kits was set at 90 d from the date of manufacture.

Final Radiopharmaceutical

Eight lots of ^{111}In -DTPA-hEGF Injection were prepared from the kits (Table 2). All radiopharmaceutical preparations met specifications for total radioactivity, pH, radiochemical purity, clarity and color, and sterility and apyrogenicity. ^{111}In -DTPA-hEGF Injection was stable for 24 h stored at 4°C. The mean radiochemical purity at 24 h was $93.1\% \pm 4.2\%$ ($n = 3$). The expiry of ^{111}In -DTPA-hEGF Injection was set at 4 h from the time of preparation.




FIGURE 3. Direct receptor-binding assay curve for binding of ^{111}In -DTPA-hEGF (prepared from kit lot 3B003) to MDA-MB-468 human breast cancer cells. K_a and B_{\max} values were 2.2×10^7 L/mol and 1.1×10^6 receptors per cell, respectively.

DISCUSSION

GMP are the foundation of a quality process that ensures that pharmaceuticals meet standards appropriate to their intended use. A central component of GMP is the establishment of specifications and standard operating procedures (SOPs) for pharmaceutical manufacturing extending from raw materials through intermediates to the final product. Guidelines for GMP have been standardized by the International Conference on Harmonization of Technical Requirements for Registration of Pharmaceuticals for Human Use (ICH) (8). Health Canada has adopted these guidelines in establishing the GMP regulations in Canada for pharmaceuticals (17) and modified them to include radiopharmaceuticals (18). In this study, we described our approach to designing GMP processes for manufacturing a kit for the preparation of ^{111}In -DTPA-hEGF Injection, a novel targeted radiotherapeutic agent for advanced EGFR-positive breast cancer. Since the processes were designed and validated in the clinical radiopharmaceutical research laboratory at the University Health Network, a University of Toronto-affiliated teaching hospital, we propose that the approach repre-

TABLE 2
Quality Control Testing of ¹¹¹In-DTPA-hEGF Injection

Test	Specification	Radiopharmaceutical lot							
		35-07/ 09/02	43-08/ 12/02	39-09/ 03/02	52-09/ 09/02	43-09/ 16/02	42-10/ 08/02	45-10/ 29/02	46-10/ 28/02
Radioactivity (MBq)	<3,083	220	151	318	151	115	168	129	132
pH	5.5–6.5	6.5	6.5	6.5	6.0	6.0	6.0	6.0	6.0
Radiochemical purity (%)	>90	94	97	97	96	99	98	96	98
Radionuclidic purity (%)	>99.9	100	100	100	100	100	100	100	100
Clarity and color	Clear and colorless	Passed	Passed	Passed	Passed	Passed	Passed	Passed	Passed
Sterility	USP XXV	Passed	Passed	Passed	Passed	Passed	Passed	Passed	Passed
Apyrogenicity	USP XXV	Passed	Passed	Not tested	Not tested	Not tested	Passed	Not tested	Not tested

sents a useful example of manufacturing an investigational biotechnology-based radiopharmaceutical in an academic setting under GMP for early phase I clinical trials in humans. The processes described are different than those of “good pharmacy practice” intended to compound radiopharmaceuticals for individual patients in a hospital setting. The unique issues in establishing GMP in a hospital setting include the low-batch sizes involved, local distribution usually only within a single institution, and the need to allocate limited resources to greatest effect. These issues were taken into consideration in designing the GMP for ¹¹¹In-DTPA-hEGF Injection.

Health Canada GMP guidelines specify that all raw materials intended for pharmaceutical use be pharmacopoeial or equivalent quality, that an identity test be performed, and that a certificate of lot analysis be obtained from the supplier confirming the purity (17). Pharmacopoeial-quality raw materials obtained for manufacturing the kits included sodium bicarbonate USP, sodium acetate USP, nitrogen NF, Sterile Water for Injection USP, and Sodium Chloride Injection USP. In addition, type 1 glass vials that met USP specifications for sterility and apyrogenicity were purchased to dispense the kits. Health Canada does not require in-house assays of materials labeled as pharmacopoeial quality (i.e., Sterile Water for Injection USP). Chloroform and DTPA dianhydride were not available in pharmacopoeial quality but were obtained in high purity (>98%). The NF assay for edetic acid was adapted to confirm the purity of DTPA dianhydride (12). Identity tests were performed on all raw materials (including those of pharmacopoeial quality) and certificates of lot analysis were obtained from the suppliers.

A major challenge in manufacturing the kits under GMP conditions was securing a source of suitable-quality hEGF. Preclinical studies of the radiopharmaceutical were conducted using hEGF_{1–53}, a high-purity (>98%) but “research-quality” material produced in *Saccharomyces cerevisiae*. This material was used to establish the specifications and analytic methods for hEGF as well as to manufacture the first 3 pilot batches of the kits. Because hEGF_{1–53} was not recommended for human use and complete details on its

production and quality control required by Health Canada were not available from the supplier, these circumstances necessitated a change in the source of hEGF raw material to hEGF_{1–51} obtained from an alternate supplier.

hEGF_{1–51} is a natural isoform of hEGF. The C-terminal amino acids Leu-52 and Arg-53 are not present in hEGF_{1–51} nor are they required for receptor binding (15). The hEGF_{1–51} material was produced in *Pichia pastoris* under ISO9001 standards (19) that are similar to GMP; therefore, the material was considered “pharmaceutical quality.” Complete manufacturing and quality control information was provided by the supplier. To demonstrate the receptor-binding equivalence of hEGF_{1–51} and hEGF_{1–53}, the ability of the 2 materials to displace the binding of ¹²³I-hEGF_{1–53} to MDA-MB-468 breast cancer cells was compared. In addition, hEGF_{1–51} was tested against specifications established for the identity and purity of hEGF. These tests showed that hEGF_{1–51} exhibited identical receptor-binding properties as hEGF_{1–53} and met or exceeded the specifications established for the raw material. Furthermore, we have recently determined that ¹¹¹In-DTPA-hEGF prepared from hEGF_{1–51} exhibits identical cytotoxic properties in vitro against MDA-MB-468 cells as that prepared from hEGF_{1–53} (unpublished data, August 2003).

It was important to create a kit formulation for preparing ¹¹¹In-DTPA-hEGF Injection because it allows rapid, simple, and reproducible preparation of the radiopharmaceutical. It also minimizes the manipulation steps involved since the very high labeling efficiency achieved (94%–99%) eliminates the need for postlabeling purification. This formulation also allows certain quality control tests (e.g., receptor-binding properties and protein purity or homogeneity) to be evaluated before patient administration and others (e.g., sterility and apyrogenicity) to be fully validated.

¹¹¹In-DTPA-hEGF exhibited specific receptor-mediated binding to MDA-MB-468 breast cancer cells (K_a , 0.9–1.1 × 10⁷ L/mol; B_{max} , 1.1–2.2 × 10⁶ sites per cell). The K_a and B_{max} values were similar to those of ¹²³I-hEGF_{1–53} (K_a , 1.6–3.4 × 10⁷ L/mol; B_{max} , 0.9–2.2 × 10⁶ sites per cell [not shown]) but the K_a values were lower than those previously

reported for ^{111}In -DTPA-hEGF (K_a , $7.5 \pm 3.8 \times 10^8 \text{ L/mol}$) (13). The B_{max} values for ^{111}In -DTPA-hEGF were similar to those previously reported (B_{max} , $1.3 \pm 0.3 \times 10^6$ sites per cell) (13). Based on the similar K_a and B_{max} values for ^{111}In -DTPA-hEGF and ^{123}I -hEGF measured using identical assay methodology, we conclude that the radiopharmaceutical exhibited preserved receptor-binding properties. There was no change in the receptor-binding properties of ^{111}In -DTPA-hEGF Injection prepared from the kits when stored for up to 90 d at 4°C.

The labeling efficiency of the kits was almost quantitative (94%–99%) when 115–318 MBq ^{111}In were added to each vial. It was further demonstrated by labeling aliquots of the kit solution with increasing amounts of ^{111}In (1.1–74 MBq) that the radiochemical purity of ^{111}In -DTPA-hEGF Injection would remain within specifications (>90%) with as much as 3,083 MBq added to each kit. Single kits were recently labeled with 740–773 MBq ^{111}In , producing ^{111}In -DTPA-hEGF Injection with a radiochemical purity of >94%. ^{111}In -DTPA-hEGF Injection was prepared by aseptically decapping the vials under laminar air flow and adding ^{111}In chloride directly into the vial using a micropipette and sterile tip. This was necessary due to the very high concentration of ^{111}In chloride radiochemical (>3.7 GBq/mL) from MDS Nordion or PerkinElmer. Since ^{111}In chloride was not pharmaceutical quality, ^{111}In -DTPA-hEGF Injection was terminally sterilized by filtration through a 0.22- μm Millex GV low-protein-binding filter. Retrospective USP Sterility and Pyrogen Tests validated the method as yielding a final product that was sterile and pyrogen free. Nevertheless, it should be possible to prepare ^{111}In -DTPA-hEGF Injection without the need for this terminal sterilization step by aseptically adding a sterile solution of ^{111}In chloride directly into the vial using a sterile syringe and needle.

In the planned phase I clinical trial, we intend to administer escalating single doses of ^{111}In -DTPA-hEGF ranging from 185 to 2,960 MBq to EGFR-positive breast cancer patients. It is anticipated therefore that 1 or 2 vials of the kit will be labeled with ^{111}In , and the corresponding administered mass of DTPA-hEGF will be 0.25–0.50 mg. ^{131}I -hEGF has been administered safely to humans for imaging squamous cell lung carcinoma in amounts up to 3.0 mg (20). Additionally, preclinical toxicology studies performed in our laboratory in mice and rabbits have shown that doses of ^{111}In -DTPA-hEGF up to 25 $\mu\text{g/kg}$ are extremely well tolerated with no evidence of significant normal tissue toxicity (unpublished data). These preclinical doses correspond to approximately 1.2–1.7 mg of ^{111}In -DTPA-hEGF in a 50- to 70-kg human.

CONCLUSION

A kit was designed and manufactured under GMP conditions for the rapid, simple, and reproducible preparation of ^{111}In -DTPA-hEGF Injection, a novel targeted radiothera-

peutic agent for advanced EGFR-positive breast cancer. Specifications, SOPs, and quality control methods were developed for all raw materials, key intermediates (pharmaceutical buffers and kits), and the final radiopharmaceutical product. ^{111}In -DTPA-hEGF Injection prepared from the kits was radiochemically pure, exhibited preserved receptor-binding properties toward EGFR-positive MDA-MB-468 human breast cancer cells, and was sterile and pyrogen free. We conclude that the kits are suitable for preparing ^{111}In -DTPA-hEGF Injection for evaluation in a planned phase I clinical trial in breast cancer patients. The GMP processes were incorporated into the Chemistry and Manufacturing section of a CTA submitted by the University Health Network to Health Canada for ^{111}In -DTPA-hEGF Injection.

ACKNOWLEDGMENT

This research was supported by grants from the U.S. Army Breast Cancer Research Program (DAMD17-02-1-0559), the Susan G. Komen Breast Cancer Foundation (BCTR0100840), and the Ontario Research and Development Challenge Fund. Parts of this study were presented at the 11th European Symposium on Radiopharmacy and Radiopharmaceuticals, Innsbruck, Austria, March 15–18, 2003.

REFERENCES

1. Mariani G, Bodei L, Adelstein SJ, Kassis A. Emerging roles for radiometabolic therapy of tumors based on Auger electron emission. *J Nucl Med.* 2000;41:1519–1521.
2. Faraggi M, Gardin I, de Labriolle-Vaylet C, Moretti J-L, Bok BD. The influence of tracer localization on the electron dose rate delivered to the cell nucleus. *J Nucl Med.* 1994;35:113–119.
3. Hofer KG, Harris CR, Smith MJ. Radiotoxicity of intracellular Ga-67, I-125 and H-3: nuclear versus cytoplasmic radiation effects in murine L1210 leukemia cells. *Int J Radiat Biol Relat Stud Phys Chem Med.* 1975;28:225–241.
4. Reilly RM, Kiarash R, Cameron RG, et al. ^{111}In -Labeled EGF is selectively radiotoxic to human breast cancer cells overexpressing EGFR. *J Nucl Med.* 2000;41:429–438.
5. Klijn JGM, Berns PMJJ, Schmitz PIM, Foekens JA. The clinical significance of epidermal growth factor receptor (EGF-R) in human breast cancer: a review on 5232 patients. *Endocr Rev.* 1992;13:3–17.
6. Chen P, Mrkobrada M, Vallis KA, et al. Comparative antiproliferative effects of novel targeted Auger electron radiotherapy, chemotherapy and external γ -radiation on EGFR-overexpressing breast cancer cells. *Nucl Med Biol.* 2002;29:693–699.
7. Chen P, Cameron R, Wang J, Vallis KA, Reilly RM. Antitumor effects and normal tissue toxicity of ^{111}In -labeled epidermal growth factor administered to athymic mice bearing EGFR-positive human breast cancer xenografts. *J Nucl Med.* 2003;44:1469–1478.
8. ICH Expert Working Group. *Good Manufacturing Practice Guide for Active Pharmaceutical Ingredients.* International Conference on Harmonization of Technical Requirements for Registration of Pharmaceuticals for Human Use. Document no. Q7A. Geneva, Switzerland: ICH Secretariat; November 10, 2000: 1–43.
9. Reagent Chemicals. *Official Specification: Chloroform.* 7th ed. Washington, DC: American Chemical Society; 1986:208–212.
10. United States Pharmacopoeia 25. *Official Monograph: Sodium Acetate.* Rockville, MD: United States Pharmacopoeial Convention; 2002:1574.
11. United States Pharmacopoeia 25. *Official Monograph: Sodium Bicarbonate.* Rockville, MD: United States Pharmacopoeial Convention; 2002:1575–1577.
12. National Formulary 20. *Official Monograph: Edetic Acid.* Rockville, MD: United States Pharmacopoeial Convention; 2002:2544.
13. Reilly RM, Kiarash R, Sandhu J, et al. A comparison of EGF and MA52 labeled with ^{111}In for imaging human breast cancer. *J Nucl Med.* 2000;41:903–911.

14. Hommel U, Harvey TS, Driscoll PC, Campbell ID. Human epidermal growth factor: high resolution solution structure and comparison with human transforming growth factor α . *J Mol Biol.* 1992;227:271–282.
15. Kuo B-S, Kusmik WF, Poole JC, et al. Pharmacokinetic evaluation of two human epidermal growth factors (hEGF51 and hEGF53) in rats. *Drug Metab Dispos.* 1992;20:23–30.
16. Prigent SA, Lemoine NR. The type 1 (EGFR-related) family of growth factor receptors and their ligands. *Prog Growth Factor Res.* 1992;4: 1–24.
17. Health Products and Food Branch Inspectorate. *Good Manufacturing Practices Guidelines.* Ottawa, Canada: Health Canada; 2002:1–86. Available at: http://www.hc-sc.gc.ca/hpfb-dgpsa/inspectorate/gmp_guidelines_2002_tc_e.html. Accessed December 31, 2002.
18. Health Products and Food Branch Inspectorate. *Annex to the GMP Guidelines: Good Manufacturing Practices for Schedule C Drugs.* Ottawa, Canada: Health Canada; 1999:1–12. Available at: http://www.hc-sc.gc.ca/hpfb-dgpsa/inspectorate/annex_sched_c_tc_e.html. Accessed June 30, 1999.
19. International Organization for Standardization. *ISO9001.* Geneva, Switzerland. 2003. Available at: <http://www.iso.ch>. Accessed December 15, 2000.
20. Cuartero-Plaza A, Martinez-Miralles E, Rosell R, Vadell-Nadal C, Farre M, Real FX. Radiolocalization of squamous lung carcinoma with ^{131}I -labeled epidermal growth factor. *Clin Cancer Res.* 1996;2:13–20.

