

REFERENCES

1. Graham MM. Ventilation–perfusion lung scanning: stuck in a rut? *J Nucl Med*. 2014;55:1395–1396.
2. Bajc M, Neilly JB, Miniati M, Schuemichen C, Meignan M, Jonson B. EANM guidelines for ventilation/perfusion scintigraphy: part 2. Algorithms and clinical considerations for diagnosis of pulmonary emboli with V/P(SPECT) and MDCT. *Eur J Nucl Med Mol Imaging*. 2009;36:1528–1538.
3. Collart JP, Roelants V, Vanpee D, et al. Is a lung perfusion scan obtained by using single photon emission computed tomography able to improve the radionuclide diagnosis of pulmonary embolism? *Nucl Med Commun*. 2002;23:1107–1113.

Michael M. Graham

University of Iowa
3863 JPP Department of Radiology
200 Hawkins Dr.
Iowa City, IA 52242

Published online Nov. 7, 2014.
DOI: 10.2967/jnumed.114.149047

Qualitative ¹⁸F-FDG PET/CT Response Evaluation After Chemotherapy or Radiotherapy for Head and Neck Squamous Cell Carcinoma: Is There an Equivocal Group?

TO THE EDITOR: We read with great interest the recent article by Marcus et al. (1). The objective of this retrospective study on 214 patients with head and neck squamous cell carcinoma was to validate qualitative interpretation criteria for ¹⁸F-FDG PET/CT assessment of response after chemoradiotherapy or radiotherapy in terms of accuracy, reader reliability, and predictive value for survival outcomes. This is an area of particular interest because the use of ¹⁸F-FDG PET/CT for response assessment of head and neck squamous cell carcinoma is becoming increasingly widespread (2). Their Hopkins criteria were used to assign a score of 1–5, with scores of 1–3 considered negative for residual disease. A score of 1 was for focal ¹⁸F-FDG uptake less than activity in the internal jugular vein, a score of 2 was for focal ¹⁸F-FDG uptake more than activity in the internal jugular vein but less than liver uptake, a score of 3 was for likely inflammatory changes, and scores of 4 and 5 were for focal uptake greater than liver uptake. The study demonstrated high interreader agreement and an overall negative predictive value of 91.1%.

The authors stated that “no established qualitative interpretation criteria...have been published” (1). However, Porceddu et al. reported final results in 2011 (3) of a high-quality prospective study on 112 patients evaluating an ¹⁸F-FDG–directed policy for the management of patients with neck node–positive head and neck squamous cell carcinoma after chemoradiotherapy or radiotherapy using qualitative PET response criteria. With some similarity to the Hopkins criteria, these response criteria were prospectively implemented; focal uptake greater than liver background was considered positive, focal uptake less than liver but more than surrounding normal tissues was considered equivocal, and no uptake above background or diffuse uptake without underlying structural abnormality was considered negative. In this prospective study, PET-based nodal assessment had a negative predictive value of 98.1% (3). The most significant difference in this method of classification is the assignment of an equivocal response to focal uptake less than liver background, which would be assigned a score of 1 or 2 and considered negative according to the Hopkins criteria.

The management of this group of equivocal responders in neck lymph nodes in an era in which neck dissections are not routinely performed (4) is a particularly difficult clinical issue. In the study by Porceddu et al. (3), 11 of 112 patients had an equivocal response and 10 of these 11 patients became negative on a repeated PET scan performed within the study protocol after a 4- to 6-wk interval and were spared a neck dissection; None of these 10 patients had subsequent neck failure. We have previously reported our initial experience with ¹⁸F-FDG PET for response assessment (5). In a recent update of our series (6), 10 of 105 patients had an equivocal response according to the reporting criteria published by Porceddu et al. (3); 2 of these 10 patients subsequently had clinicopathologic evidence of lymph node disease.

We believe that the clinical significance and optimal management of focal ¹⁸F-FDG PET uptake below the level of liver background remains uncertain. In light of these differing qualitative response criteria and the higher negative predictive value reported by Porceddu et al. (3) for a negative PET scan, it is possible that the negative predictive value provided by the Hopkins criteria may be improved by separately considering patients with an equivocal response as defined by Porceddu et al. (3). We would be interested to learn whether there is any difference in the negative predictive value reported by Marcus et al. (1) comparing patients who scored a 1, 2, or 3 according to the Hopkins criteria and classified as having a negative scan.

REFERENCES

1. Marcus C, Ciarallo A, Tahari AK, et al. Head and neck PET/CT: therapy response interpretation criteria (Hopkins criteria)—interreader reliability, accuracy, and survival outcomes. *J Nucl Med*. 2014;55:1411–1416.
2. Gupta T, Master Z, Kannan S, et al. Diagnostic performance of post-treatment FDG PET or FDG PET/CT imaging in head and neck cancer: a systematic review and meta-analysis. *Eur J Nucl Med Mol Imaging*. 2011;38:2083–2095.
3. Porceddu SV, Pryor DI, Burnmeister E, et al. Results of a prospective study of positron emission tomography-directed management of residual nodal abnormalities in node-positive head and neck cancer after definitive radiotherapy with or without systemic therapy. *Head Neck*. 2011;33:1675–1682.
4. Ferlito A, Corry J, Silver CE, Shaha AR, Thomas Robbins K, Rinaldo A. Planned neck dissection for patients with complete response to chemoradiotherapy: a concept approaching obsolescence. *Head Neck*. 2010;32:253–261.
5. Prestwich RJ, Subesinghe M, Gilbert A, Chowdhury FU, Sen M, Scarsbrook AF. Delayed response assessment with FDG-PET-CT following (chemo)radiotherapy for locally advanced head and neck squamous cell carcinoma. *Clin Radiol*. 2012;67:966–975.
6. Slevin F, Ramasamy S, Sen M, Subesinghe M, Scarsbrook A, Prestwich R. Delayed response assessment FDG PET-CT following (chemo)radiotherapy for head and neck squamous cell carcinoma [abstract]. *Radiother Oncol*. 2014;111(suppl 1):266.

Robin Prestwich

Mehmet Sen

Andrew Scarsbrook*

*St. James Institute of Oncology

Level 1, Bexley Wing

Leeds LS9 7TF, U.K.

E-mail: a.scarsbrook@nhs.net

Published online Nov. 26, 2014.
DOI: 10.2967/jnumed.114.148114

REPLY: We agree that having an equivocal group is a challenging clinical issue in assessing patients with head and neck squamous cell carcinoma after chemoradiation therapy. The Hopkins criteria (1) are a simple, standardized, qualitative method of assessing therapy,