

MAX S. LIN

Veterans Administration Hospital
and Stanford University School of
Medicine, Palo Alto, California

GERALD BURKE

Michael Reese Hospital and Medical Center
Chicago, Illinois

REFERENCES

1. HOSAIN F, MCINTYRE PA, POULOSE K, et al: Binding

of trace amounts of ionic indium-113m to plasma transferrin. *Clin Chim Acta* 24: 69-75, 1969

2. WOCHNER RD, ADATEPE M, VAN AMBURG A, et al: A new method for estimation of plasma volume with the use of the distribution space of indium-113m-transferrin. *J Lab Clin Med* 75: 711-720, 1970

3. STRAUSS HW, MCINTYRE PA, WAGNER HN: Evaluation of indium-labeled transferrin for the determination of plasma volume. *J Nucl Med* 10: 442, 1969

4. LIN MS, BURKE G, LEE JI: The variability of early urinary excretion of ^{113m}In. *J Nucl Med* 14: 126-127, 1973

SUPERIOR VENA CAVAL OBSTRUCTION AND INCREASED RADIOCOLLOID

ACTIVITY ON LIVER SCINTIPHOTOS

Recent case reports in the *Journal of Nuclear Medicine* have recorded the unusual occurrence of focal increases in radiocolloid activity on liver scintiphotos (1-3). We have recently observed two additional such cases. Both patients presented with superior vena caval obstruction secondary to metastatic carcinoma. In each case static and dynamic scintiphotos demonstrated focal areas of increased colloid at the junction of the right and left hepatic lobes (Fig. 1).

Four of five previously reported cases of radiocolloid "hot spots" have also been associated with metastatic carcinoma and superior vena caval obstruction. The fifth case was due to a hemangioma (4). Caval portal shunting due to vena caval obstruction would appear a likely mechanism for this phenomenon (2). Metastatic tumors do not contain Kupffer's cells and "colloid-concentrating" metastatic tumors have yet to be documented.

G. BRUCE HOPKINS

Scripps Memorial Hospital
La Jolla, California

REFERENCES

1. COEL M, HALPERN SE, ALAZRAKI NP, et al: Intra-hepatic lesion presenting as an area of increased radiocolloid uptake on a liver scan. *J Nucl Med* 13: 221-222, 1972

2. HOLMQUIST DL, BURDINE JA: Caval-portal shunting as a cause of a focal increase in radiocolloid uptake in normal livers. *J Nucl Med* 14: 348-351, 1973

3. MRKOLAJKOW A, JASINSKI WK: Increased focal uptake of radiocolloid by the liver. *J Nucl Med* 14: 175, 1973

4. VOLPE JA, JOHNSTON GS: "Hot" hepatic hemangioma: A unique radiocolloid-concentrating liver scan lesion. *J Surg Oncology* 2: 373-377, 1970




FIG. 1. Dynamic (1) and static (2) scintiphotos of radiocolloid hot spot. Patient had superior vena caval obstruction secondary to metastatic lung carcinoma.

ERRATA

In the Works in Progress abstract entitled "¹¹¹In-dium-Bleomycin: A New Radiopharmaceutical for Tumor Scintigraphy" (*J Nucl Med* 14: 641, 1973) Melvin J. Silverstein should be added to the list of authors. The correct sequence of authors is: Ramesh C. Verma, Leslie R. Bennett, Juan J. Touya, Melvin J. Silverstein, Donald L. Morton, and Ewa Witt.

The Letter to the Editor by N. Adishesan (*J Nucl Med* 14: 722, 1973) should be entitled "Radiostrontium Localization in Normal Lungs?" and the Author's Reply (*J Nucl Med* 14: 723, 1973) by T. K. Chaudhuri should be entitled "Radiostrontium Deposition in Lungs: 'Apparently Normal' Lungs versus Occult Aspergillosis.'