

out by Rivera, its constant position and decreasing activity with time should help avoid any erroneous interpretation of the study.

I am delighted to hear of this observation with the use of a long-available radiopharmaceutical that I believe is underutilized. I am optimistic that the future availability of  $^{128}\text{I}$ -rose bengal will rekindle

interest in this stimulating and useful diagnostic study in jaundiced patients.

LEONARD M. FREEMAN

Albert Einstein College of Medicine  
Bronx, New York

## CANCER OF THE COLON VISUALIZED BY STRONTIUM SCINTIGRAPHY

With regard to a paper by Chaudhuri, et al (1) concerning a case in which there was scintigraphic ( $^{87\text{m}}\text{Sr}$ ) demonstration of liver metastases secondary to a primary carcinoma of the colon, allow me to make the following comments.

The scintigraphic demonstration of reactivated colorectal cancer and especially the demonstration of local recurrences is of considerable importance for the radiotherapeutic management of these patients. As in most of the cases, an abdominoperineal resection has been performed, barium enema and digital or bimanual examinations are not feasible, and consequently the identification and localization of a recurrence is difficult until it acquires a large size. Excretory urography is not always contributory whereas other more sophisticated procedures such as arteriography and venography are time-consuming and cause considerable discomfort to the patients. Scintigraphy is a simple and easy-to-perform method but unfortunately tumors do not always concentrate radionuclides. Chaudhuri, et al (1) postulate a special affinity of strontium to cancer cells, but if this were the case one would expect more frequent visualization of colonic cancers, either primary or metastatic. We routinely investigate colorectal cancer patients with scintigraphy using various radionuclides. We have employed  $^{67}\text{Ga}$  and bleomycin labeled with  $^{67}\text{Co}$ , but better results have been obtained with  $^{87\text{m}}\text{Sr}$ . Radioactive strontium has been useful mainly in demonstrating: (A) bone metastases and (B) recurrences in the minor pelvis, either intraosseous or extraosseous. A few cases of extraosseous recurrences in the pelvis demonstrated by  $^{87\text{m}}\text{Sr}$  have already been reported (2,3). Although it may be true that strontium concentrates in extraosseous tumors according to any one of the mechanisms reported by Samuels (4) and Chaudhuri (1), it also may be true that these mechanisms differ from case to case. I am inclined to believe, however, that in the majority of cases, strontium localizes in colonic cancer because it mimics calcium. This assumption

is based on the following facts: (A) it is known that primary mucinous colonic adenocarcinoma occasionally undergoes calcification often detected radiologically; (B) recurrences in the minor pelvis may become necrotic due to inadequate vascular supply. In these necrotic regions, calcium may be deposited. Then too, previous radiation therapy facilitates calcification. (C) It has been repeatedly reported that liver metastases from primary cancer of the colon may present hazy or stippled calcifications, often detectable—but not always—on the radiographs. As a matter of fact if such a calcification is recognized, the possibility of colonic cancer should be considered (5). (D) In the case of a recurrence in the minor pelvis, Denonvillier's fascia and the bladder wall constitute a barrier to the forward spread of the tumor. The tumor, therefore, extends posteriorly to the presacral space and the sacrum. Periosteal reaction of this bone may result in increased radioactivity although there is no actual bone involvement.

CONSTANTINE PAPAVALIOU

Alexandra University Hospital  
Athens, Greece

## REFERENCES

1. CHAUDHURI TK, CHAUDHURI TK, GO RT, et al: Uptake of  $^{87\text{m}}\text{Sr}$  by liver metastasis from Ca of the colon. *J Nucl Med* 14: 295-296, 1973
2. PAPAVALIOU C, KOSTAMIS P, ANGELAKIS P, et al: Localization of  $^{87\text{m}}\text{Sr}$  in extraosseous tumors. *J Nucl Med* 12: 265-268, 1971
3. PAPAVALIOU C, KOSTAMIS P, CONSTANTINIDES C, et al: La valeur de la scintigraphie au  $^{87\text{m}}\text{Sr}$  dans la radiothérapie palliative du cancer rectocolique récidivant. *J Radiol Electrol Med Nucl* 52: 497-500, 1971
4. SAMUELS LD: Detection and localization of extra-skeletal malignant neoplasms of children with strontium  $^{87\text{m}}$ . *Am J Roentgenol Radium Ther Nucl Med* 115: 777-782, 1972
5. FRED HL, EINBAND JM, COLLINS LC: Calcifications in intra-abdominal and retroperitoneal metastases. *Am J Roentgenol Radium Ther Nucl Med* 91: 138-148, 1964