

as an alternative explanation for "hot emboli," the possible association with hypercoagulability of blood, as seen in critically ill cancer patients and in postoperative patients.

In the present case, the latter two explanations are not applicable; rather, the radioactive microemboli appear to have arisen from tagged deep venous thromboses following low-pressure, low-volume injections soon after a contrast venogram had been performed. It appears, therefore, that radioactive microembolization arising from a tagged thrombus may occur as a complication of radionuclide venography in a patient with an abnormal peripheral venous system.

D. M. CONCA
D. R. BRILL
J. D. SHOOP
Geisinger Medical Center
Danville, Pennsylvania

REFERENCES

1. LUTZKER LG, PEREZ LA: Radioactive embolization from upper-extremity thrombophlebitis. *J Nucl Med* 16: 241-242, 1975
2. KONTZEN FN, BESCHI RJ, PITTMAN JA JR: Radioactive pulmonary emboli. *Ala J Med Sci* 8: 27-29, 1971
3. PRESTON DF, GREENLAW RH: "Hot spots" in lung scans. *J Nucl Med* 11: 422-425, 1970

Diethyl-IDA: A Promising Hepatobiliary Radiopharmaceutical

The recent editorial on hepatobiliary radiopharmaceuticals has come to our attention (1). Indeed, diethyl-IDA, used for several months for hepatobiliary studies in our department, appears to be the best tracer available, especially in disorders of the gallbladder. With diethyl-IDA, neither false negative nor false positive results were ever observed, while this was certainly not the case with 99m-Tc-PyG or 99m-Tc-DHTA.

We recently reviewed a series of 30 patients investigated with diethyl-IDA, comprising 20 normal subjects and 10 patients suffering from various gallbladder disorders (8 cholecystitis and 2 cholelithiasis). Eight of 20 normals were not fasting at the moment of the investigation. In 18 cases out of the 20 normal subjects, the gallbladder was visualized from 9 to 34 min after injection of the tracer. In two other normals, the gallbladder was not seen, but none of these was fasting. Failure to visualize the gallbladder was systematically observed in all the pathologic cases, even when late views were obtained.

Thus, provided that the investigation was performed in the fasting state, a normal gallbladder was always visualized; but any acute or subacute disorder of the gallbladder was characterized by the absence of visualization of the gallbladder.

S. PAUWELS
M. STEELS
L. PIRET
C. BECKERS
University of Louvain
Brussels, Belgium

REFERENCE

1. RONAI PM: Hepatobiliary radiopharmaceuticals: Defining their clinical role will be a galling experience. *J Nucl Med* 18: 488-490, 1977

Pericardial Accumulation of Tc-99m Methylene Diphosphonate in a Case of Pericarditis

Technetium-99m phosphate tracers are currently being employed for the detection of acute myocardial infarcts (1-3). While imaging for myocardial infarct, we observed an unusual case of accumulation of Tc-99m methylene diphosphonate (MDP) in the pericardium of a patient with pericarditis and a small pericardial effusion following acute myocardial infarction.

A 32-year-old white male was admitted to the hospital with acute onset of crushing retrosternal chest pain accompanied by nausea and diaphoresis. An electrocardiogram revealed typical changes of an acute infero-lateral wall infarction. This was further confirmed by the characteristic rise in cardiac enzymes, with a peak CPK of 350 (normal, less than 12), peak SGOT of 561 (normal, less than 40), and a peak LDH of 1230 (normal, less than 250). Serial electrocardiograms revealed typical evolution of an infero-lateral infarction with development of Q waves in I, II, III, AVL, AVF, V5, and V6.

Twenty-four hours after admission the patient complained of precordial pain suggestive of pericarditis—worse on inspiration and on lying on the left side, and relieved slightly by leaning forward. A triphasic friction rub was audible at this time. Chest x-ray showed mild venous congestion and interstitial edema. Heart size was normal. On the third day of infarction, myocardial scintigraphy was performed using 15 mCi Tc-99m MDP, accumulating 500,000 counts per image. The images demonstrated uptake compatible with an infero-lateral myocardial infarction, but there was also unusual accumulation of the tracer in the pericardial silhouette (Fig. 1). An echocardiogram done immediately following the scan revealed a small (5-mm) pericardial effusion. Coronary arteriography revealed single-vessel disease involving the left circumflex artery at the origin of the first marginal branch. An 80% stenosis was estimated, with good distal runoff. Left ventriculography revealed marked hypokinesis of the anterior, inferior and apical surfaces. Hemodynamic values were normal. BUN, creatinine, serum calcium and phosphorus were normal. Repeat chest x-ray did not show any calcification within the pericardial silhouette.

The mechanism of the phenomenon of soft-tissue uptake is not well understood. Both benign and malignant soft-tissue lesions may concentrate phosphate tracers (4). Recently, intense myocardial uptake of Tc-99m diphosphonate has been reported in a uremic patient with secondary hyperparathyroidism and pericarditis (5). Concentration of these tracers in inflammatory conditions (6,7) and pleural effusions (8) have also been described. Accumulation of Tc-99m MDP in our case is a very rare condition. Since the pericardial effusion in our patient was quite small, it is likely that the increased uptake was due to the inflammatory component of the pericarditis rather than to the effusion, although direct proof of this is lacking.

AMOLAK SINGH
MICHAEL USHER
LAWRENCE RAPHAEL
Jewish General Hospital and McGill University
Montreal, Quebec, Canada