

potheses and experiments designed to test them, whose goal is to show an effect of the caloric content of a test meal on gastric emptying. Because of this, I found the discussion of the nice study by Phillips and his colleagues of special interest (1).

The caloric content of a meal is measured by placing it in a calorimeter, oxidizing it completely, and measuring the heat produced. A living organism has no physiologic mechanism to measure a meal's calories, much less react to them. In fact, the calories of a meal are released over a long period of time after it is digested—probably days. An acute effect of an ingested meal's calories on gastric motility, or anything else for that matter, is preposterous.

The problem presented to investigators of this bankrupt idea is that it is impossible to prepare two meals that only differ in caloric content and do not differ in many already recognized and physiologically important variables. In summary, I find calories influencing gastric motility indigestible.

## REFERENCES

1. Phillips WT, Schwartz JG, Blumhardt R, McMahan CA. Linear gastric emptying of hyperosmolar glucose solutions. *J Nucl Med* 1991;32:377-381.

**Robert S. Hattner**

*University of California, San Francisco  
San Francisco, California*

**REPLY:** We agree with Dr. Hattner that gastric emptying is determined by multiple complex factors. The meal phase (1) (liquid versus solid), composition of the meal (2) (carbohydrate:protein:fat), weight (2) and volume (3) of the meal, stimulation of osmoreceptors (4), as well as energy density (5,6) (kcal/g) have all been reported to affect the rate of gastric emptying.

Although many of the articles in nuclear medicine and gastric motility literature emphasize the effect of the caloric content of a test meal on gastric emptying (5-7) (an increased number of calories is associated with slower gastric emptying), this theory is not always consistent. Our article (*J Nucl Med* 1991;32:377-381) described a 400-kcal glucose solution emptying almost twice as slowly as our 200-kcal glucose solution. However, we described in the same article a 400-kcal liquid glucose solution having a longer half-emptying time than the 600-kcal meal in the Velchik et al. study (107 min with the 400-kcal glucose solution versus 95 min with the 600-kcal solid meal). This difference, we believe, may be due to a slower rate of intestinal absorption of the solid meal, compared to the more rapid intestinal absorption of the glucose solution.

Thus, although we believe that all the variables mentioned above affect gastric emptying, in our opinion it is the rate of absorption of nutrients by the small intestine that is the most important factor controlling gastric emptying. The release of at least one gut hormone, gastric inhibitory polypeptide (GIP), has been shown to be dependent on the rate of absorption of nutrients by the small intestine (8). Another possibility is that circulating blood levels of nutrients could exert a strong control on gastric emptying. Several investigations have shown that intravenous infusion of glucose greatly slows gastric emptying (9,10). Obviously, the rate of absorption of nutrients by the small intestine and circulating blood levels are closely connected.

This rate of absorption, as Dr. Hattner has mentioned, can

vary greatly between different types of foods (11). The absorption rate is dependent on the physical properties of each food.

The use of glucose solutions, as in our studies, allows for very rapid intestinal absorption after the glucose is emptied from the stomach. In this case, the gastric emptying time should be increasingly prolonged with increasing caloric content, caloric content being simply an indirect way of specifying the quantity of glucose.

If these ideas are correct, they have important applications in the performance of gastric emptying studies. Therefore, we believe that, in order to have reproducible gastric emptying studies, the physical properties and quantity (weight) of a gastric emptying meal must be rigidly controlled.

## REFERENCES

1. Heading RC, Tothill P, McLoughin GP, Shearman DJC. Gastric emptying rate measurement in man. *Gastroenterology* 1976;71:45-50.
2. Moore JG, Christian PE, Coleman RE. Gastric emptying of varying meal weight and composition in man. *Dig Dis Sci* 1981;26:16-22.
3. Malmud LS, Fisher RS, Knight LC, Rock E. Scintigraphic evaluation of gastric emptying. *Semin Nucl Med* 1982;12:116-125.
4. Elias E, Gibson GJ, Greenwood LF, Hunt JN, Tripp JH. The slowing of gastric emptying by monosaccharides and disaccharides in test meals. *J Physiol* 1968;194:317-326.
5. Velchik MG, Reynolds JC, Alavi A. The effect of meal energy content on gastric emptying. *J Nucl Med* 1989;30:1106-1110.
6. McHugh PR, Moran TH. Calories and gastric regulatory capacity with implications for feeding. *Am J Physiol* 1979;236:R254-R260.
7. Brener W, Hendrix TR, McHugh PR. Regulation of the gastric emptying of glucose. *Gastroenterology* 1983;85:76-82.
8. Cretzfeldt W, Ebert R. New developments in the incretin concept. *Diabetologia* 1985;28:565-573.
9. Morgan LM, Tredger JAT, Hampton SM, French AP, Peake JCF, Marks V. The effect of dietary modification and hyperglycaemia on gastric emptying and gastric inhibitory polypeptide (GIP) secretion. *Br J Nutr* 1988;60:29-37.
10. MacGregor IL, Gueller R, Watts HD, Meyer JH. The effect of acute hyperglycemia on gastric emptying in man. *Gastroenterology* 1976;70:190-196.
11. Wolever TMS. The glycemic index. In: Bourne GH, ed. *Aspects of some vitamins, minerals and enzymes in health and disease*. World Rev Nutr Diet. Basel: Karger, 1990:120-185.

**W. T. Phillips**

**J. G. Schwartz**

**R. Blumhardt**

**C. Alex McMahan**

*The University of Texas*

*Health Science Center at San Antonio*

*San Antonio, Texas*

## A Novel In Vivo Red Cell-Specific Label

**TO THE EDITOR:** In his editorial (1), Ronald J. Callahan listed 13 radionuclides that have been used to label human red blood cells and commented on the in vitro and in vivo labeling protocols. We would like to add radioiodine to the list following our work on the selective labeling of human erythrocytes by para-iodobenzenesulphonamide (pIBS). The results of these studies have recently been published (2-4), and we simply wish to point out the advantages of our technique over current methodologies.

Our targeting method is based on the use of a facilely radioiodinated, lipophilic enzyme-inhibitor as a radionuclide "carrier." Red cells are ideal for this approach since they contain a high concentration of the target enzyme, carbonic anhydrase. Further-