
The Journal of Nuclear Medicine

JNMM

Volume 34, Number 2 • February 1993

173 Evaluating Paragangliomas

180 Imaging Procedures for Pheochromocytomas

222 Technetium-99m-Tetrofosmin

281 Gated and Ungated SPECT Technetium-99m-MIBI

288 FDG-PET in Bone Tumor

A Full Table of Contents Begins on Page 4A, Annotations on Pages 7A-8A



The Official Publication of
The Society of Nuclear Medicine, Inc.

Iodine-131-MIBG Scintigraphy in Preoperative and Postoperative Evaluation of Paragangliomas: Comparison with CT and MRI Imaging

Iodine-131-MIBG scintigraphy, CT and MRI were used to evaluate 36 patients with clinically suspect functioning paragangliomas.....page 173

Editorial: Can We Agree on the Best Imaging Procedure(s) for Localization of Pheochromocytomas?

.....page 180

Splenectomy in Idiopathic Thrombocytopenic Purpura: Its Correlation with the Sequestration of Autologous Indium-111-Platelets

A retrospective analysis of ¹¹¹In-platelet sequestration in 111 patients diagnosed with idiopathic thrombocytopenic purpura is provided.page 182

Assessment of Myocardial Perfusion Defect Size After Early and Delayed SPECT Imaging with Technetium-99m-Hexakis 2-Methoxyisobutyl After Stress

The assumption that ^{99m}Tc-MIBI undergoes minimal change during the first four hours postinjection was tested in 20 patients with coronary artery disease by SPECT studies after exercise stress or dipyridamole infusion.page 187

Left Ventricular Cavity-to-Myocardium Count Ratio: A New Parameter for Detecting Resting Left Ventricular Dysfunction Directly from Tomographic Thallium Perfusion Scintigraphy

A retrospective study of 80 patients who were undergoing evaluation of chest discomfort, or with known or suspected coronary disease, was used to evaluate a quantitative index which relates thallium activity in the left ventricular cavity to that in the myocardium.....page 193

Noninvasive, Real-Time Monitoring of Renal Function: The Ambulatory Renal Monitor

A radiation detector attached to a miniature data logger was used to monitor the clearance of a glomerular filtration agent from the extracellular space in 50 patientspage 199

The Dosimetry of Iodine-123-Labeled TISCH: A SPECT Imaging Agent for the D1 Dopamine Receptor

Dose to seven healthy subjects was estimated by the absorbed fraction technique. Conjugate images of the body were serially acquired for up to 24 hr. Count rates in visualized organs were measured on each image.page 208

Hepatobiliary Scintigraphy in a Pediatric Population: Determination of Hepatic Extraction Fraction by Deconvolution Analysis

To assess the feasibility of measuring hepatic extraction fraction and hepatic clearance half-times in pediatric patients, 45 children in four categories were studied: normal (12), obstruction (9), hepatocellular disease (16) and miscellaneous (8).page 214

Technetium-99m-Tetrofosmin as a New Radiopharmaceutical for Myocardial Perfusion

A new cationic complex of ^{99m}Tc has been synthesized and evaluated for use in myocardial perfusion imaging.page 222

Potential Gallium-68 Tracers for Imaging the Heart with PET: Evaluation of Four Gallium Complexes with Functionalized Tripodal Tris(Salicylaldimine) Ligands

Four new tripodal hexadentate tris(salicylaldimine) ligands are evaluated and assessed for their potential in providing novel ⁶⁸Ga radiopharmaceuticalspage 228

Rapid Infarct Imaging with a Technetium-99m-Labeled

Antimyosin Recombinant Single-Chain Fv: Evaluation in a Canine Model of Acute Myocardial Infarction

In vivo targeting of acute myocardial infarction with a novel recombinant single-chain Fv antimyosin protein was attempted in the belief that the use of smaller labeled fragments may result in improved image quality.page 234

Effect of Hypoxia on the Uptake of Technetium-99m-Glucuronate and Technetium-99m-Gluconate by Chinese Hamster Ovary Cells In Vitro

The accumulation of ^{99m}Tc-glucuronate in zones of acute cerebral injury and tumors was compared with ^{99m}Tc-gluconate in an in vitro system of cultured Chinese hamster ovary fibroblastspage 242

Synthesis and PET Studies of Fluorine-18-BMY 14802: A Potential Antipsychotic Drug

Fluorine-18-labeled BMY 14802 was synthesized and resolved to obtain enantiomerically pure samples and its behavior was examined in a baboon with PET.page 246

Uptake of Technetium-99m-Tebroxime in Cultured Myocardial Cells: Comparison with Thallium-201 and Technetium-99m-Sestamibi

The authors assessed the effect of mild to strong metabolic inhibition on the uptake of tebroxime in comparison with the uptake of ²⁰¹Tl and sestamibi.page 255

Rhenium-186-Labeled Monoclonal Antibodies for Radioimmunotherapy: Preparation and Evaluation

A simple technique for labeling IgM, IgG, fragmented antibodies and tumor necrosis factor (TNF) with ¹⁸⁶Re is presented.page 260

Formulation of Technetium-99m-Aerosol Colloid with Improved Delivery Efficiency for Lung

Ventilation Imaging

An in vitro system was used to compare the delivery of various ^{99m}Tc radiopharmaceuticals from nebulizer solutions. The addition of a phosphate buffer produced a promising tin colloid based agent. *page 268*

A Dual-Isotope Comparison of the Distribution of Radioiodinated-E-17 α -Iodovinyl-11 β -Methoxyestradiol and 1-Iodo-1, 1-Bis(4-Hydroxyphenyl)-Phenylethylene Estrogens in the Immature Female Rat

The specific uptake, retention and distribution in immature rat tissue of non-steroidal and steroidal estrogen were compared in order to assess their potential for therapy and/or diagnosis of ER-containing cancers in patients. *page 272*

Inhomogeneity of Gated and Ungated SPECT Technetium-99m-Sestamibi Bull's-Eyes in Normal Dogs: Comparison with Thallium-201

Simulation predictions for both ECG-gated and ungated SPECT sestamibi and ^{201}Tl myocardial perfusion images, established previously by computer model, were validated by studies of four normal dogs. *page 281*

Fluorine-18-Fluorodeoxyglucose Positron Emission Tomography in Technetium-99m-Hydroxymethylenediphosphate Negative Bone Tumors

Two cases of bone tumors with high FDG uptake and negative ^{99m}Tc -HMDP bone scintigraphy suggested a complementary role for FDG-PET in the detection and follow-up of bone tumors. *page 288*

Cerebral Blood Flow and Magnetic Resonance Imaging in Locked-in Syndrome

The cerebral blood flow of a patient suffering from locked-in syndrome

was examined before and after onset using ^{99m}Tc -hexamethylpropyleneamine oxime SPECT and intravenous ^{133}Xe *page 291*

Technetium-99m-Perchnetate Scintigraphy in an Ileal Duplication of the Stomach and Duodenum

A case of ileal duplication of the stomach and duodenum was found in an 8-mo-old hispanic male presenting with gastrointestinal bleeding. *page 294*

Scintigraphic "Doughnut Sign" on Skeletal Imaging due to a Hemangioendothelioma of Bone

Bone scans of a 61-yr-old patient revealed multiple lesions of the tibia and foot, several of which were doughnut-shaped and corresponded to lytic abnormalities on radiographs. *page 297*

Results of Surgery for Portacaval Shunt in a Patient with Situs Inversus Evaluated by Per Rectal Portal Scintigraphy

The authors treated a 70-yr-old patient presenting with situs inversus, asplenia and a large portacaval shunt and observed the clinical course by per rectal portal scintigraphy. *page 301*

A Quantitative Assessment of Patient Motion and Its Effect on Myocardial Perfusion SPECT Images

The effects of patient motion on ^{201}Tl SPECT scintigrams were assessed with computer simulation to create 66 image sets with artifactual vertical, horizontal and combined patient motion. *page 303*

Three-Dimensional Registration of Cardiac Positron Emission Tomography Attenuation Scans

A method for three-dimensional registration of two cardiac PET attenuation scans of the same subject acquired at different times was evaluated in 15

subjects. *page 311*

Automated Detection of the Intercommissural Line for Stereotactic Localization of Functional Brain Images

Seven subjects participated in validation studies of a new technique for automated detection of the intercommissural line for PET. *page 322*

The Channel Ratio Method of Scatter Correction for Radionuclide Image Quantitation

The change in the ratio of counts from two symmetrical adjacent energy windows straddling the photo peak was used to eliminate the contribution of scattering photons during imaging with ^{99m}Tc *page 330*

Automated Region Definition for Cardiac Nitrogen-13-Ammonia PET Imaging

Six volunteers were studied with ^{13}N -ammonia PET as part of an effort to develop an analytical method for the quantification of regional myocardial blood flow in the clinical setting. *page 338*

Quality Assurance of White Blood Cell Labeling with a Test Based on Adherence

A quality control assay was developed on the premise that proper labeling should not affect the adherence characteristics of white blood cells to nylon fibers. *page 345*

The Present State of Landfill Disposal of Radioactive Materials

The problems hospitals may face when private corporations make policies for the disposal of low-level radioactive waste that are more stringent than existing laws are discussed. *page 349*