

2. Osbakken MD, Boucher GA, Okada RD, Bingham JB, Strauss HW, Pohost GM. Spectrum of global left ventricular responses to supine exercise: limitation in the use of ejection fraction in identifying patients with coronary artery disease. *Am J Cardiol* 1983;51:28-35.
3. Crawford MH, White DH, Amon KW. Echocardiography evaluation of left ventricular size and performance during handgrip and supine and upright bicycle exercise. *Circulation* 1979;59:1188-1196.
4. Gorlin R, Cohen LS, Elliott WC, Klein MD, Lane FJ. Effect of supine exercise on left ventricular volume and oxygen consumption in man. *Circulation* 1965;32:361-371.
5. Slutsky R, Karliner J, Ricci D, Schuler G, Pfisterer M, Peterson K, Ashburn W. Response of left ventricular volume to exercise in man assessed by radionuclide equilibrium angiography. *Circulation* 1979;60:565-571.
6. Tamaki N, Gill JB, Moore RH, Yasuda T, Boucher CA, Strauss HW. Cardiac responses to daily activities and exercise in normal subjects assessed by an ambulatory ventricular function monitor. *Am J Cardiol* 1987;59:1164-1169.
7. Kiess MC, Dimsdale JE, Moore RH, Liu P, Newell J. The effects of stress on left ventricular ejection fraction. *Eur J Nucl Med* 1988;14:12-16.
8. Koike A, Haruki I, Koichi T, Hiroe M. Detecting abnormalities in left ventricular function during exercise by respiratory measurement. *Circulation* 1989;80:1736-1746.
9. Boucher CA, Anderson M, Schneider MS, Murphy J, Okada R. Left ventricular function before and after reaching the anaerobic threshold. *Chest* 1985;87:145-150.
10. Rodrigues A, Maddahi J, Brown H, et al. Responses of left and right ventricular ejection fractions to aerobic and anaerobic phases of upright and supine exercise in normal subjects. *Am Heart J* 1989;118:319-324.
11. Pace L, Cuocolo A, Nappi A, Nicolai E, Trimarco B, Salvatore M. Accuracy and repeatability of left ventricular systolic and diastolic function measurements using an ambulatory radionuclide monitor. *Eur J Nucl Med* 1992;19:800-806.
12. Cuocolo A, Nappi A, Nicolai E, et al. Ambulatory monitoring of left ventricular function: evaluation of transient left ventricular dysfunction in patients with coronary artery disease. *G Ital Cardiol* 1992;22:813-824.
13. Jones NL. *Clinical exercise testing*. Philadelphia: WB Saunders; 1988:208-212.
14. Pace L, Cuocolo A, Mangoni di S, et al. Left ventricular systolic and diastolic function measurements using an ambulatory radionuclide monitor: effects of different time averaging on accuracy. *J Nucl Med* 1993;34:1602-1606.
15. Weber KT, Janicki SJ. Cardiopulmonary exercise testing for evaluation of chronic cardiac failure. *Am J Cardiol* 1985;55:22A-31A.
16. Poliner LR, Deheimer GJ, Lewis SE, Parkey RW, Blomquist CG, Willerson JT. Left ventricular performance in normal subjects: a comparison of the response to exercise in the upright and supine position. *Circulation* 1980;62:528-534.
17. Breisblatt WM, Weiland FL, McLain JR, Tomlinson GC, Burns MJ, Spaccavento LJ. Usefulness of ambulatory radionuclide monitoring of left ventricular function early after acute myocardial infarction for predicting residual myocardial ischemia. *Am J Cardiol* 1988;62:1005-1010.
18. Matsumura N, Nishijima H, Koizumi S, Hashimoto F, Yasuda H. Determination of anaerobic threshold for assessment of functional state in patients with chronic heart failure. *Circulation* 1983;68:360-367.
19. Opasich C, Cobelli F, Riccardi G, Aquilani R, Specchia G. Applicability of the anaerobic threshold in patients with previous myocardial infarction. *Adv Cardiol* 1986;35:100-107.

ERRATUM

Due to a production error, in the December 1994 issue of *JNM*, the radiotracer on page 1922 in the article "Dopamine D₂ Receptor Imaging with Iodine-123-Iodobenzamide SPECT in Idiopathic Rotational Torticollis" by Hierholzer et al., was misprinted. The correct radiotracer is Iodine-123-IBZM.