

outflow resistance, resulting in decreased interventricular septal oxygen demand (3).

A similar hypothesis was considered by Ono et al. (2) who wrote "...there is the possibility that oxygen demand in the septum is decreased because of reduced thickening..." but they dismissed this on the basis of increased intramyocardial pressure during the phase of the cardiac cycle incorporating the major component of left anterior descending flow ("...suggesting that an asynchronous but reserved myocardial contraction was occurring). In their Figure 5, simultaneous coronary flow and intramyocardial pressure tracings are displayed without registration to the electrocardiogram. During pacing-induced left bundle branch block, two phenomena are illustrated. First, mean intramyocardial pressure rises, consistent with tachycardia-induced decrease in left ventricular dimensions and increase in myocardial thickness and compression (4). Second, phasic increase in intramyocardial pressure in the septum is more synchronous with the major component of left anterior descending coronary flow. It is consistent with measured thallium uptake that this component of coronary flow is predominantly to nonseptal left ventricular segments. That septal intramyocardial pressure is dramatically "out of phase" with this flow is explicable if the QRS complex duration is long (e.g., 150 msec.) in relation to the paced R-R interval (e.g., 300 msec.). This would result in sufficient ventricular asynchrony to cause a large "phase shift" in intramyocardial pressure of the septum. In this way, left anterior descending flow is maximal during relaxation of the left ventricular free wall and contraction of the septum. That the septum may in effect function more as a wall of the right rather than the left ventricle and have a lower metabolic demand is thereby well illustrated by this work.

I accept the important findings that coronary flow to and metabolic activity of the intraventricular septum are reduced in left bundle branch block without evidence of ischemia. Perhaps diminished glucose uptake does in fact reflect decreased septal metabolic demand as the proximate cause of its diminished perfusion in left bundle branch block. How this might affect the utility of glucose uptake to evaluate the viability of hibernating and other myocardial segments with asynchronous activation is an intriguing subject for conjecture and further investigation.

## REFERENCES


1. Altehoefer C, vom Dahl J, Bares R, Stocklin GL, Bull U. Metabolic mismatch of septal beta-oxidation and glucose utilization in left bundle branch block assessed with PET. *J Nucl Med* 1995;36:2056-2059.
2. Ono S, Nohara R, Kambara H, Okuda K, Kawai C. Regional myocardial perfusion and glucose metabolism in experimental left bundle branch block. *Circulation* 1992;85:1125-1131.
3. Burns RJ, Galligan L, Wright LM, Lawand S, Burke R, Gladstone PJ. Improved specificity of myocardial thallium-201 single-photon emission computed tomography in patients with left bundle branch block by dipyridamole. *Am J Cardiol* 1991;68:504-508.
4. Wigger CL. The interplay of coronary vascular resistance and myocardial compression in regulating coronary flow. *Circ Res* 1954;2:271-278.

R.J. Burns  
University of Toronto  
Toronto, Ontario, Canada

## Double-Phase Technetium-99m-Sestamibi Scanning to Evaluate Nodular Thyroid Malignancy

**TO THE EDITOR:** In a recent article, Vattimo et al. (1) reported on double-phase thyroid scintigraphy with <sup>99m</sup>Tc-MIBI in patients with cold solitary nodules (1). They found that the technique could identify patients with Hürthle cell tumors showing persistent uptake on the late image (3-4 hr postinjection). In contrast, papillary and follicular malignant nodules showed complete washout. Moreover, they concluded that persistent MIBI uptake seems to be characteristic of the Hürthle cell tumors.

In our own experience, double-phase scanning of the thyroid gland with <sup>99m</sup>Tc-MIBI at 2 hr is a useful test to evaluate single thyroid nodules. The nodular retention of the radiotracer on delayed images at 2 hr postinjection



**FIGURE 1.** Two-hour delayed MIBI image reveals nodular retention of the radiotracer in malignant papillary (A) and follicular lesions (B).

is an indicator of malignancy regardless the histological type of the lesion (Fig. 1).

Vattimo et al. also stated that Földes et al. (2) performed the double-phase technique with results similar to theirs. These authors, however, acquired delayed images at 1 hr postinjection in 10 of 58 patients, including many with multinodular goiter.

Although more experience is needed to evaluate the diagnostic accuracy of the test, we believe that nodular washout of radiotracer in malignant papillary and follicular nodules is not observed if late images are acquired at 2 hr postinjection.

## REFERENCES

1. Vattimo A, Bertelli P, Cintonio M, Burroni L, Volterrani D, Vella A. Identification of Hürthle cell tumor by single-injection, double-phase scintigraphy with technetium-99m-sestamibi. *J Nucl Med* 1995;36:778-782.
2. Földes I, Lévy A, Stotz G. Comparative scanning of thyroid nodules with technetium-99m-pertechnetate and technetium-99m-methoxyisobutylisonitrite. *Eur J Nucl Med* 1993;20:330-333.

Omar Alonso  
Fernando Mut  
Graciela Lago

Clinical Hospital of the University of Uruguay  
Montevideo, Uruguay

**REPLY:** We thank Alonso et al. for their interest in our article (1). We observed that Hürthle cell tumors (HCT) take up and retain <sup>99m</sup>Tc-sestamibi (MIBI), in contrast with other thyroid tumors that take up and release the tracer. Therefore, we believe that single-injection, double-phase (15-30 min and 3-4 hr p.i.) scintigraphy with MIBI can detect cold thyroid nodules bearing HCT. MIBI uptake by thyroid nodules is related to vascularity and cellularity (2), but its retention is mainly related to mitochondrial concentration (3) and secondarily to the initial uptake: As a result, visual interpretation could be misleading since the retention could be either true or apparent according to the washout rate. For this reason, we calculated the washout rate and the nodule-to-thyroid uptake ratio (N/T). We observed HCTs that exhibited a slow washout rate (<20% h<sup>-1</sup>) and increased N/T in comparison with other thyroid tumors presenting high washout rates (>20% h<sup>-1</sup>) and decreased N/T. This scintigraphic pattern is related to a histopathological feature of such tumors (high density of oxyphilic cells due to crowded mitochondria).

We are not surprised by the observations of Alonso et al., since a similar case has already been reported by Taillefer et al. (4). We also observed cases of non-HCTs exhibiting delayed retention of MIBI (adenoma and follicular carcinoma). In such patients, the N/T decreased and the washout rate was high (>20% h<sup>-1</sup>) so that their retention was quite apparent (Fig. 1). However, we are indebted to Alonso et al. for their observation, which we will consider in the evaluation of future patients.