

outflow resistance, resulting in decreased interventricular septal oxygen demand (3).

A similar hypothesis was considered by Ono et al. (2) who wrote "...there is the possibility that oxygen demand in the septum is decreased because of reduced thickening..." but they dismissed this on the basis of increased intramyocardial pressure during the phase of the cardiac cycle incorporating the major component of left anterior descending flow ("...suggesting that an asynchronous but reserved myocardial contraction was occurring). In their Figure 5, simultaneous coronary flow and intramyocardial pressure tracings are displayed without registration to the electrocardiogram. During pacing-induced left bundle branch block, two phenomena are illustrated. First, mean intramyocardial pressure rises, consistent with tachycardia-induced decrease in left ventricular dimensions and increase in myocardial thickness and compression (4). Second, phasic increase in intramyocardial pressure in the septum is more synchronous with the major component of left anterior descending coronary flow. It is consistent with measured thallium uptake that this component of coronary flow is predominantly to nonseptal left ventricular segments. That septal intramyocardial pressure is dramatically "out of phase" with this flow is explicable if the QRS complex duration is long (e.g., 150 msec.) in relation to the paced R-R interval (e.g., 300 msec.). This would result in sufficient ventricular asynchrony to cause a large "phase shift" in intramyocardial pressure of the septum. In this way, left anterior descending flow is maximal during relaxation of the left ventricular free wall and contraction of the septum. That the septum may in effect function more as a wall of the right rather than the left ventricle and have a lower metabolic demand is thereby well illustrated by this work.

I accept the important findings that coronary flow to and metabolic activity of the intraventricular septum are reduced in left bundle branch block without evidence of ischemia. Perhaps diminished glucose uptake does in fact reflect decreased septal metabolic demand as the proximate cause of its diminished perfusion in left bundle branch block. How this might affect the utility of glucose uptake to evaluate the viability of hibernating and other myocardial segments with asynchronous activation is an intriguing subject for conjecture and further investigation.

REFERENCES

1. Altehoefer C, vom Dahl J, Bares R, Stocklin GL, Bull U. Metabolic mismatch of septal beta-oxidation and glucose utilization in left bundle branch block assessed with PET. *J Nucl Med* 1995;36:2056-2059.
2. Ono S, Nohara R, Kambara H, Okuda K, Kawai C. Regional myocardial perfusion and glucose metabolism in experimental left bundle branch block. *Circulation* 1992;85:1125-1131.
3. Burns RJ, Galligan L, Wright LM, Lawand S, Burke R, Gladstone PJ. Improved specificity of myocardial thallium-201 single-photon emission computed tomography in patients with left bundle branch block by dipyridamole. *Am J Cardiol* 1991;68:504-508.
4. Wigger CL. The interplay of coronary vascular resistance and myocardial compression in regulating coronary flow. *Circ Res* 1954;2:271-278.

R.J. Burns

University of Toronto
Toronto, Ontario, Canada

Double-Phase Technetium-99m-Sestamibi Scanning to Evaluate Nodular Thyroid Malignancy

TO THE EDITOR: In a recent article, Vattimo et al. (1) reported on double-phase thyroid scintigraphy with ^{99m}Tc-MIBI in patients with cold solitary nodules (1). They found that the technique could identify patients with Hürthle cell tumors showing persistent uptake on the late image (3-4 hr postinjection). In contrast, papillary and follicular malignant nodules showed complete washout. Moreover, they concluded that persistent MIBI uptake seems to be characteristic of the Hürthle cell tumors.

In our own experience, double-phase scanning of the thyroid gland with ^{99m}Tc-MIBI at 2 hr is a useful test to evaluate single thyroid nodules. The nodular retention of the radiotracer on delayed images at 2 hr postinjection




FIGURE 1. Two-hour delayed MIBI image reveals nodular retention of the radiotracer in malignant papillary (A) and follicular lesions (B).

is an indicator of malignancy regardless the histological type of the lesion (Fig. 1).

Vattimo et al. also stated that Földes et al. (2) performed the double-phase technique with results similar to theirs. These authors, however, acquired delayed images at 1 hr postinjection in 10 of 58 patients, including many with multinodular goiter.

Although more experience is needed to evaluate the diagnostic accuracy of the test, we believe that nodular washout of radiotracer in malignant papillary and follicular nodules is not observed if late images are acquired at 2 hr postinjection.

REFERENCES

1. Vattimo A, Bertelli P, Cintorino M, Burroni L, Volterrani D, Vella A. Identification of Hürthle cell tumor by single-injection, double-phase scintigraphy with technetium-99m-sestamibi. *J Nucl Med* 1995;36:778-782.
2. Földes I, Lévy A, Stotz G. Comparative scanning of thyroid nodules with technetium-99m-pertechnetate and technetium-99m-methoxyisobutylisonitrile. *Eur J Nucl Med* 1993;20:330-333.

Omar Alonso

Fernando Mut

Graciela Lago

Clinical Hospital of the University of Uruguay
Montevideo, Uruguay

REPLY: We thank Alonso et al. for their interest in our article (1). We observed that Hürthle cell tumors (HCT) take up and retain ^{99m}Tc-sestamibi (MIBI), in contrast with other thyroid tumors that take up and release the tracer. Therefore, we believe that single-injection, double-phase (15-30 min and 3-4 hr p.i.) scintigraphy with MIBI can detect cold thyroid nodules bearing HCT. MIBI uptake by thyroid nodules is related to vascularity and cellularity (2), but its retention is mainly related to mitochondrial concentration (3) and secondarily to the initial uptake: As a result, visual interpretation could be misleading since the retention could be either true or apparent according to the washout rate. For this reason, we calculated the washout rate and the nodule-to-thyroid uptake ratio (N/T). We observed HCTs that exhibited a slow washout rate (<20% h⁻¹) and increased N/T in comparison with other thyroid tumors presenting high washout rates (>20% h⁻¹) and decreased N/T. This scintigraphic pattern is related to a histopathological feature of such tumors (high density of oxyphilic cells due to crowded mitochondria).

We are not surprised by the observations of Alonso et al., since a similar case has already been reported by Taillefer et al. (4). We also observed cases of non-HCTs exhibiting delayed retention of MIBI (adenoma and follicular carcinoma). In such patients, the N/T decreased and the washout rate was high (>20% h⁻¹) so that their retention was quite apparent (Fig. 1). However, we are indebted to Alonso et al. for their observation, which we will consider in the evaluation of future patients.




FIGURE 1. Cold thyroid nodules (left column) show intense MIBI uptake in the early image (center). Intense retention in Patient A ($WO = 9\% h^{-1}$) and faint retention in Patient B ($WO = 24\% h^{-1}$; right column). Histopathology revealed HCT in Patient A and follicular carcinoma in Patient B.

REFERENCES

1. Vattimo A, Bertelli P, Cintorino M, et al. Identification of Hürthle cell tumor by single-injection, double-phase scintigraphy with technetium-99m-sestamibi. *J Nucl Med* 1995;36:778–782.
2. Vattimo A, Bertelli P, Burrioni L. Effective visualization of suppressed thyroid tissue by means of baseline methoxy-isobutyl-isonitrile in comparison with ^{99m}Tc -pertechnetate scintigraphy after TSH stimulation. *J Nucl Biol Med* 1992;36:315–318.
3. Carvalho PA, Chiu ML, Kronauge JF, et al. Subcellular distribution and analysis of technetium-99m-MIBI in isolated perfused rat hearts. *J Nucl Med* 1992;33:1516–1531.
4. Taillefer R, Boucher Y, Potvin C, et al. Detection and localization of parathyroid adenoma in patients with hyperparathyroidism using a single radionuclide imaging procedure with technetium-99m-sestamibi (double-phase study). *J Nucl Med* 1992;23:1801–1807.

Angelo Vattimo
Paolo Bertelli
Marcella Cintorino
Luca Burrioni
Duccio Volterrani
Alessandra Vella
University of Siena
Siena, Italy

(continued from page 1823)

FIRST IMPRESSIONS Hepatic Uptake of Bone Tracer




Figure 1.




Figure 2.




Figure 3.

PURPOSE

A 56-yr-old man presenting with back pain was referred for bone scanning. In the initial bone scan, both kidneys were visualized with some uneven activity. A well-defined photopenic area was observed on the lateral side of the right kidney (Fig. 1). Additional activity was also observed cephalad to the right kidney at the 10-11 costal area. Tumor in the right kidney with the possibility of bone metastases, accessory kidney or tumor in the soft tissue were considered. The patient underwent surgery and histology verified a clear-cell carcinoma in the right kidney. Postoperatively, the symptoms were reduced, but activity in the costal region remained unchanged and appeared to be mobile (Fig. 2). Ultrasound and CT demonstrated the presence of "gallbladder" in this area (Fig. 3). There was no evidence of bone metastases. Apparently, ^{99m}Tc -MDP was retained in the gallbladder. Possible mechanisms of hepatic bone tracer uptake include decreased pH of the eluate used

in preparing the MDP or excess aluminum ion concentration. This explanation is not entirely satisfactory since this patient's gallbladder is visualized on both the preoperative and postoperative bone scans.

TRACER

Technetium-99m-MDP

ROUTE OF ADMINISTRATION

Intravenous injection

TIME AFTER INJECTION

Three hours

INSTRUMENTATION

Philips gamma camera

CONTRIBUTORS

Bob Dugal and Øivind O. Ing, Central Hospital, Fredrikstad, Norway.