

ment of Veterans Affairs. Thanks also are ended to Robin Sawyer for her assistance in the preparation of this manuscript.

REFERENCES

1. Dalen JE, Alpert JS. Natural history of pulmonary embolism. *Prog Cardiovasc Diseases* 1975;17:259-270.
2. Goldhaber SZ, Hennekens CH, Evans DA, Newton EC, Godleski JJ. Factors associated with correct antemortem diagnosis of major pulmonary embolism. *Am J Med* 1982;73:822-826.
3. Stein PD, Willis PW, DeMets DL. History and physical examination in acute pulmonary embolism in patients without preexisting cardiac or pulmonary disease. *Am J Cardiol* 1981;45:218-223.
4. Hoellerich VL, Wigton RS. Diagnosing pulmonary embolism using clinical findings. *Arch Intern Med* 1986;146:1699-1704.
5. Stein PD, Terrin ML, Hales CA, et al. Clinical, laboratory, roentgenographic and electrocardiographic findings in patients with acute pulmonary embolism and no pre-existing cardiac or pulmonary disease. *Chest* 1991;100:598-603.
6. Hoffman JM, Lee A, Grafton ST, et al. Clinical signs and symptoms in pulmonary embolism—a reassessment. *Clin Nucl Med* 1994;19:803-808.
7. McNeil BJ. Ventilation-perfusion studies and the diagnosis of pulmonary embolism: concise communication. *J Nucl Med* 1980;21:319-323.
8. Braun SD, Newman GE, Ford K, et al. Ventilation-perfusion scanning and pulmonary angiography: correlation in clinical high-probability pulmonary embolism. *Am J Roentgenol* 1984;143:977-980.
9. The PLOPED Investigators. Value of the ventilation/perfusion scan in acute pulmonary embolism—results of the prospective investigation of pulmonary embolism diagnosis (PIOPED). *JAMA* 1990;263:2753-2759.
10. Kelley MA, Carson JL, Palevsky HI, Schwartz JS. Diagnosing pulmonary embolism: new facts and strategies. *Ann Int Med* 1991;114:300-306.
11. Davis RB, Schauwecker DS, Siddiqui AR, et al. Indeterminate lung imaging—can the number be reduced? *Clin Nucl Med* 1986;11:577-582.
12. Quinn RJ, Butler SP. A decision analysis approach to the treatment of patients with suspected pulmonary emboli and an intermediate probability lung scan. *J Nucl Med* 1991;32:2050-2056.
13. Smith R, Maher JM, Miller RI, Alderson PO. Clinical outcomes of patients with suspected pulmonary embolism and low-probability aerosol-perfusion scintigrams. *Radiology* 1987;164:731-733.
14. Lee ME, Biello DR, Kumar B, Siegel BA. "Low-probability" ventilation-perfusion scintigrams: clinical outcomes in 99 patients. *Radiology* 1985;156:497-500.
15. Kahn D, Bushnell DL, Dean R, Perlman SB. Clinical outcome of patients with a "low probability" of pulmonary embolism on ventilation-perfusion lung scan. *Arch Intern Med* 1989;149:377-379.
16. Oudkerk M, van Beek EJR, van Putten WLJ, Buller HR. Cost-effectiveness analysis of various strategies in the diagnostic management of pulmonary embolism. *Arch Intern Med* 1993;153:947-954.
17. Sullivan DC, Coleman RE, Mills SR, Ravin CE, Hedlund LW. Lung scan interpretation: effect of different observers and different criteria. *Radiology* 1983;149:803-807.
18. Quinn RJ, Nour R, Butler SP, et al. Pulmonary embolism in patients with intermediate probability lung scans: diagnosis with doppler venous US and D-Dimer measurement. *Radiology* 1994;190:509-511.
19. Saenger EL, Buncher CR, Specker BL, McDevitt RA. Determination of clinical efficacy: nuclear medicine as applied to lung scanning. *J Nucl Med* 1985;26:793-806.
20. Frankel N, Coleman RE, Prior DB, Sostman HD, Ravin CE. Utilization of lung scans by clinicians. *J Nucl Med* 1986;27:366-369.
21. Biello DR, Mattar AG, McKnight RC, et al. Ventilation-perfusion studies in suspected pulmonary embolism. *Am J Roentgenol* 1979;133:1033-1037.
22. Biello DR, Mattar AG, Osei-Wusu A, et al. Interpretation of indeterminate lung scans. *Radiology* 1979;133:189-194.
23. Freitas JE, Sarosi MG, Nagle CC, et al. Modified PLOPED criteria used in clinical practice. *J Nucl Med* 1995;36:1573-1578.
24. Catania TA, Caride VJ. Single perfusion defect and pulmonary embolism: angiographic correlation. *J Nucl Med* 1990;31:296-301.
25. Spies WG, Burstein SP, Dillehay GL, et al. Ventilation-perfusion scintigraphy in suspected pulmonary embolism: correlation with pulmonary angiography and refinement of criteria for interpretation. *Radiology* 1986;159:383-390.
26. Gottschalk A, Sostman HD, Coleman RE, et al. Ventilation-perfusion scintigraphy in the PLOPED study. Part II. Evaluation of the scintigraphic criteria and interpretations. *J Nucl Med* 1993;34:1119-1126.
27. Dismuke SE, Wagner EH. Pulmonary embolism as a cause of death—the changing mortality in hospitalized patients. *JAMA* 1986;255:2039-2042.
28. Kruit WHJ, De Boer AC, Sing AK, Van Roon F. The significance of venography in the management of patients with clinically suspected pulmonary embolism. *J Intern Med* 1991;230:333-339.
29. Dalen JE. When can treatment be withheld in patients with suspected pulmonary embolism? *Arch Intern Med* 1993;153:1415-1418.
30. Stein PD, Athanasoulis C, Alavi A, et al. Complications and validity of pulmonary angiography in acute pulmonary embolism. *Circulation* 1992;85:462-468.
31. Elgazzar AH, Silberstein EB, Hughes J. Perfusion and ventilation scans in patients with extensive obstructive airway disease: utility of single-breath (washin) xenon-133. *J Nucl Med* 1995;36:64-67.

Scintigraphic Evaluation of Experimental Colitis in Rabbits

Wim J.G. Oyen, Otto C. Boerman, Els T.M. Dams, Gert Storm, Louis van Bloois, Emile B. Koenders, Urbain J.G.M. van Haelst, Jos W.M. van der Meer and Frans H.M. Corstens

University Hospital Nijmegen, Departments of Nuclear Medicine, Internal Medicine and Pathology, Nijmegen; and Utrecht University, Institute for Pharmaceutical Science (UIPS), Department of Pharmaceutics, Utrecht, The Netherlands

Scintigraphic techniques are frequently used for evaluation of inflammatory bowel disease. The radiopharmaceutical of choice is labeled leukocytes. In this study, two new agents, ^{111}In -labeled polyethylene glycol-coated liposomes and ^{111}In -labeled human nonspecific gamma globulin (immunoglobulin G; IgG), were compared with ^{111}In -leukocytes in a rabbit model of colitis. **Methods:** In rabbits, acute colitis was induced by colonic instillation of trinitrobenzene sulfonic acid at 25 cm from the anal sphincter. After 24 hr, 15 MBq of the radiopharmaceuticals was injected intravenously in groups of four rabbits. Twenty-four hours after injection, the animals were killed and macroscopic abnormalities were scored in seven consecutive affected colonic segments of 5 cm each (0 = normal, 1 = inflammation, 2 = ulcers). The ex vivo uptake was measured in the normal ascending colon and the affected colonic segments. The colitis index (CI, affected-to-normal colon-uptake ratio) was calculated. **Results:** Histologically, an acute, patchy, transmural colitis was observed at the site of instillation and the distal colon. The CI of all agents in colitis lesions correlated with the severity of the abnormalities. With increasing severity, the CI for liposomes was

1.86 ± 0.24 , 4.88 ± 0.42 and 7.42 ± 0.54 ($r^2 = 0.68$, $p < 0.001$); for leukocytes 1.77 ± 0.32 , 3.10 ± 0.58 and 5.54 ± 0.83 ($r^2 = 0.31$, $p < 0.01$); for IgG 1.60 ± 0.29 , 2.81 ± 0.21 and 2.65 ± 0.21 ($r^2 = 0.29$, $p < 0.02$). **Conclusion:** Indium-111-labeled-leukocytes, -IgG and -liposomes all show increased uptake in inflamed colonic tissue. Indium-111-liposomes showed the highest CI, which correlates best with the morphological abnormalities. Indium-111-leukocytes and ^{111}In -liposomes are superior to ^{111}In -IgG for this indication.

Key Words: radionuclide imaging; diagnostic imaging; gamma globulin; indium-111; scintigraphy; inflammatory bowel disease

J Nucl Med 1997; 38:1596-1600

Inflammatory bowel disease is a condition with fluctuating episodes of relapses and remissions of acute colitis. In clinical practice, diagnostic procedures are most helpful for evaluating the status of the diseased colon. The major diagnostic tools are endoscopy (allowing direct inspection of the diseased mucosa), radiographic evaluation using barium enemas (providing images of the morphological abnormalities) and scintigraphic modalities (showing functional images of the degree of inflammatory activity in affected areas in the gut). For the latter technique, a variety of radiopharmaceuticals are available.

Received Jun. 21, 1996; revision accepted Dec. 10, 1996.

For correspondence or reprints contact: Wim J.G. Oyen, MD, University Hospital Nijmegen, Department of Nuclear Medicine, P.O. Box 9101, 6500 HB Nijmegen, The Netherlands.

When scintigraphic evaluation of the abdomen is considered labeled leukocytes are currently the agent of choice, since high sensitivity and specificity is observed, especially in acute colitis (1). Unfortunately, the preparation of radiolabeled autologous leukocytes has several disadvantages that limit its application. In particular, the necessity to draw and handle possibly contaminated blood constitutes an inherent risk to other patients and staff (2,3). Moreover, isolation and labeling of leukocytes is a relatively complicated and time-consuming procedure that is not possible in every nuclear medicine facility (4). To overcome these difficulties, research has been directed toward developing agents that are as equally effective, but easier to produce from instant, ready-to-use radiolabeling kits.

In this study in a rabbit model of acute colitis, the targeting of ^{111}In -labeled-leukocytes is compared with the performance of two recently developed radiopharmaceuticals: ^{111}In -labeled human nonspecific polyclonal gamma globulin (immunoglobulin G; IgG) and ^{111}In -labeled sterically stabilized liposomes. Indium-111-IgG has already been applied in clinical practice for the diagnosis of a wide variety of infectious and inflammatory processes (5). Labeled liposomes have also been investigated in patients (6,7). However, these conventional liposomes, as used in the past, are rapidly cleared from the circulation by phagocytic cells of the mononuclear phagocyte system (8). The advantage of sterically stabilized liposomes is the prolonged residence time in circulation with consequently higher levels of uptake in inflammatory foci and low liver and spleen uptake (9-11).

MATERIALS AND METHODS

Animal Model

In female New Zealand white rabbits (weight 2.5-3 kg), acute colitis was induced as described previously with minor modifications (12-14). Experiments were performed in accordance with the local animal welfare committee guidelines. During the experiment, the rabbits were fasted, but had water ad libitum. Animals were anesthetized with an intravenous injection of a 0.7-ml mixture of fentanyl 0.315 mg/ml and fluanisone 10 mg/ml (Hypnorm). After retrograde insertion of a flexible silicone tube, 1 ml of 30 mg trinitrobenzene sulfonic acid in 30% ethanol, followed by 1 ml 50% ethanol flush, was instilled in the colon 25 cm from the anal sphincter. Thereafter, anesthesia was terminated by intravenous injection of 0.2 ml of naloxon hydrochloride 0.4 mg/ml (Narcan). Twenty-four hours after colitis induction, the respective radiopharmaceuticals were injected through the ear vein.

Radiopharmaceuticals

Indium-111-Leukocytes. One hundred milliliters of blood were drawn carefully from an anesthetized donor rabbit by carotic artery cannulation in 60 ml syringes, each containing 10 ml 0.33% methylcellulose in citric acid dextrose. The total leukocyte count of the donor rabbit was 10×10^9 /liter, with approximately 50% granulocytes. Preparation of the labeled leukocytes was performed as described previously (15,16). In brief, after sedimentation of the erythrocytes, the supernatant was removed. The remaining cell suspension was centrifuged twice and the cell pellet was washed with phosphate buffered saline (PBS)/1% HSA. Indium-111-oxine (100 MBq) in 0.2 M tris(hydroxymethyl)aminomethane hydrochloride (pH = 8.0) was added to the cell suspension. After a 30-min incubation period, the cells were incubated at room temperature for 30 min, centrifuged and resuspended in PBS/1% human serum albumin. Morphological integrity of the leukocytes was checked by light microscopic examination. Labeling efficiency (cell associated activity/total activity $\times 100\%$) was higher than 95%. Indium-111-leukocytes (15 MBq) were administered intravenously.

Indium-111-Liposomes. Partially hydrogenated egg-phosphatidylcholine with an iodine value of 40 was derived (17). The polyethylene glycol 1900 derivative of distearoyl phosphatidylethanolamine was prepared as described previously (18). A chloroform/methanol mixture (10/1, v/v) containing polyethylene glycol-distearoyl phosphatidylethanolamine, partially hydrogenated egg-phosphatidylcholine and cholesterol was prepared in a molar ratio of 0.15:1.85:1. A lipid film was formed by rotary evaporation followed by high vacuum to remove residual organic solvent (19). The lipids were dispersed at room temperature in 6 mM Desferal in 0.9% HEPES buffer (10 mM HEPES/135 mM sodium chloride, pH 7.5) at an initial phospholipid concentration of 120 mM. The liposomes were sequentially extruded through polycarbonate filters of 0.2, 0.1, 0.08 and 0.05- μm pore size. Untrapped Desferal was removed by gel filtration on an EconoPac 10 DG column. The particle size distribution was determined by dynamic light scattering with a Malvern 4700 system using a 25 mW Helium-Neon laser. The data were analyzed using the Automeasure 3.2 software. As a measure of particle size distribution of the dispersion, a polydispersity index was determined, ranging from 0.0 (entirely monodisperse dispersion) to 1.0 (completely polydisperse dispersion). The mean size of the liposome dispersions was 90 nm with a polydispersity index of approximately 0.1.

Preformed Desferal-containing liposomes were labeled with ^{111}In essentially as described previously (10,20). Indium-111 was transported over the lipid bilayer in the form of ^{111}In -oxine and trapped irreversibly in the internal aqueous phase by the encapsulated Desferal. Briefly, the liposomes (45 mmol phospholipid/ml) were incubated for 30 min at room temperature with 200 kBq ^{111}In -oxine per millimole of phospholipid. Removal of unencapsulated ^{111}In -oxine was achieved by gel filtration on a 10DG Econo Pak column. More than 95% of the radiolabel was associated with the liposomes. Indium-111-liposomes (15 MBq) were injected intravenously.

Indium-111-IgG. Human, nonspecific polyclonal IgG was conjugated to diethylenetriaminepentaacetic bicyclic anhydride (bicyclic DTPA) as described by Hnatowich et al. and labeled with ^{111}In -chloride (21). Labeling efficiency as determined by instant thin-layer chromatography was higher than 95%. Indium-111-IgG (15 MBq) was injected intravenously.

Histology

Two rabbits were killed with an overdose of sodium phenobarbital 24 hr after colitis induction. The colon was resected and tissues were fixed in 4% formaldehyde in PBS. Tissue samples were embedded in paraffin and 5- μm sections were stained with hematoxylin-eosin for light-microscopic examination.

Imaging Procedure and Biodistribution

Twenty-four hours after colitis induction, three groups of four rabbits were intravenously injected through the ear vein with either ^{111}In -leukocytes, ^{111}In -liposomes or ^{111}In -IgG. The rabbits were immobilized in a mold and placed prone on a gamma camera equipped with a parallel-hole medium-energy collimator. Each rabbit was imaged at 5 min, 1 hr, 4 hr, 10 hr and 20 hr after injection. Images (100,000 counts per rabbit) were obtained and stored in a 256×256 matrix.

After acquiring the final images, the rabbits were killed with an overdose of sodium phenobarbital and biodistribution of the radiopharmaceuticals was determined. Blood was obtained by cardiac puncture. Tissues [muscle, lung, liver, spleen, kidney; normal (ascending) colon and affected (transverse and descending) colon] were dissected. The affected colon was divided into seven consecutive affected colonic segments of 5 cm each. The colonic segments were scored macroscopically on an arbitrary scale (0 = normal, 1 = inflammation, 2 = ulcers). The samples were weighed and their radioactivity was measured in a shielded well scintillation




FIGURE 1. Histology of trinitrobenzene sulfonic acid-induced acute colitis. The mucosa shows beginning necrosis and ulceration (arrows). The submucosa is edematous (*). Areas of granulocyte infiltration can be identified throughout the colonic wall. Hematoxylin-eosin-stained transverse sections: (A) $\times 130$, (B) $\times 330$.

gamma counter. To correct for physical decay and to permit calculation of the uptake of the radiopharmaceuticals in each organ as a fraction of the injected dose, aliquots of the injected dose were counted simultaneously. The results were expressed as percent injected dose per gram (%ID/g). From these results, the colitis index (CI, affected-to-normal colon ratio) for each of the radiopharmaceuticals was calculated in all segments.

Statistical Analysis

All mean values are given as %ID/g or ratios \pm one s.e.m. Statistical analysis was performed using one way analysis of variance. Correlations were calculated by linear regression analysis. The level of significance was set at $p < 0.05$.

RESULTS

Histologically, multiple areas of mucosal necrosis were observed at the site of instillation and the part of the colon between instillation site and rectum. The submucosa was edematous and transmural granulocytic infiltration was seen (Fig. 1).

Over time, increasing accumulation of the radiopharmaceuticals in the diseased colon was observed on the scintigraphic images. Figure 2 shows images of rabbits recorded 10 and 20 hr after injection of the radiopharmaceuticals. Physiological uptake in kidneys (liposomes and IgG only), liver and spleen was seen. Furthermore, affected parts of the colon were delineated with all radiopharmaceuticals.

The biodistribution in tissue samples is given in Table 1. Indium-111-leukocyte blood levels were lower than those of ^{111}In -IgG and ^{111}In -liposomes. This faster clearance of ^{111}In -leukocytes was also reflected in significantly lower uptake in




FIGURE 2. Scintigraphic images of rabbits with experimental colitis. (A) Indium-111 leukocytes, 10 hr postinjection. (B) Indium-111 leukocytes, 20 hr postinjection. (C) Indium-111 liposomes, 10 hr postinjection. (D) Indium-111 liposomes, 20 hr postinjection. (E) Indium-111 IgG, 10 hr postinjection. (F) Indium-111 IgG, 20 hr postinjection. Note the physiological uptake in liver, spleen and for ^{111}In -liposomes and ^{111}In -IgG also kidney and blood pool. \Rightarrow S = liver; * = spleen; \rightarrow = delineation of colitis.

muscle, lung and uptake in nonaffected normal colon. The uptake in liver and spleen markedly differed between the various preparations. Indium-111-leukocytes had much higher splenic and—to a lesser extent—liver uptake than ^{111}In -liposomes and ^{111}In -IgG. Overall absolute uptake in the affected colon was significantly higher for ^{111}In -liposomes than for the two other agents. Absolute uptake of ^{111}In -leukocytes was relatively low. Still, as indicated below, abnormalities were also well delineated with ^{111}In -leukocytes, since uptake in nonaffected colon was extremely low and consequently CIs relatively high.

CIs (diseased-to-normal colon ratios) are given in Figure 3. When comparing the relative uptake in macroscopically normal colon (scored as Grade 0), the CIs for all three preparations were similar. In nonulcerative inflammation (scored as Grade 1) and ulceration (scored as Grade 2), the relative uptake of ^{111}In -liposomes in the diseased segments was higher than that of both ^{111}In -leukocytes and ^{111}In -IgG, reflected in significantly higher CIs of ^{111}In -liposomes ($p < 0.05$ and $p < 0.001$, respectively). The relative uptake of ^{111}In -IgG was significantly lower than that of ^{111}In -leukocytes in ulceration (Grade 2) ($p < 0.05$).

When evaluating the relative uptake in the affected segments for the individual radiopharmaceuticals, a positive correlation was observed between CIs and the severity of the macroscopic abnormalities. For ^{111}In -liposomes, correlation was strongest ($r^2 = 0.68$, $p < 0.001$). Similarly, for ^{111}In -leukocytes and ^{111}In -IgG, a correlation was observed but less striking ($r^2 = 0.31$, $p < 0.01$, and $r^2 = 0.29$, $p < 0.02$, respectively).

DISCUSSION

In this study, ^{111}In -labeled sterically stabilized liposomes were shown to be a superior imaging agent for evaluation of

TABLE 1

Tissue Distribution (%ID/g; mean \pm s.e.m.) Obtained 48 hr after Colitis Induction and 24 hr after Injection of the Radiopharmaceuticals

Tissue	$^{111}\text{In-IgG}$	p^*	$^{111}\text{In-leukocytes}$	p^\dagger	$^{111}\text{In-liposomes}$
Blood	0.22 \pm 0.03	< 0.05	0.13 \pm 0.008	< 0.05	0.22 \pm 0.01
Muscle	0.01 \pm 0.001	< 0.01	0.002 \pm 0.0002	< 0.01	0.01 \pm 0.001
Lung	0.09 \pm 0.001	< 0.01	0.06 \pm 0.002	< 0.01	0.10 \pm 0.001
Liver	0.20 \pm 0.02	< 0.01	0.53 \pm 0.04	< 0.01	0.15 \pm 0.03
Spleen	0.12 \pm 0.008	< 0.01	8.08 \pm 1.09	< 0.01	0.42 \pm 0.04 [‡]
Kidney	0.19 \pm 0.02	< 0.01	0.05 \pm 0.006	< 0.01	0.20 \pm 0.02
Ascending colon (normal)	0.05 \pm 0.002	< 0.01	0.01 \pm 0.003	< 0.01	0.06 \pm 0.007
Descending colon (affected, 35 cm)	0.13 \pm 0.01	< 0.01	0.03 \pm 0.006	< 0.01	0.37 \pm 0.003 [‡]

IgG = gamma globulin.

^{*} $^{111}\text{In-IgG}$ vs. $^{111}\text{In-leukocytes}$.[†] $^{111}\text{In-leukocytes}$ vs. $^{111}\text{In-liposomes}$.[‡] $^{111}\text{In-liposomes}$ vs. $^{111}\text{In-IgG}$ ($p < 0.01$).

acute trinitrobenzene sulfonic acid-induced colitis in rabbits. In clinical practice, leukocytes are considered the standard technique for scintigraphic evaluation of disease activity in inflammatory bowel disease (1,22). In this study, $^{111}\text{In-liposomes}$ showed higher absolute, as well as uptake, ratios in diseased colonic segments compared with the $^{111}\text{In-leukocytes}$. Moreover, correlation of the macroscopic abnormalities observed in the affected colon and relative colonic uptake, of $^{111}\text{In-liposomes}$ was better compared with $^{111}\text{In-leukocytes}$. In addition, hepatosplenic uptake of $^{111}\text{In-liposomes}$ was considerably lower than that of $^{111}\text{In-leukocytes}$, which is not only important for adequate scintigraphic evaluation of the alimentary tract in the upper abdomen, but also for reduction of the radiation burden to the spleen and to long-living T-lymphocytes, which are also labeled in mixed leukocyte preparations (23). Despite the lower absolute uptake of $^{111}\text{In-leukocytes}$ in colitis lesions, diseased colon segments could be delineated well since uptake in the background tissues (i.e., normal colon and muscle) was extremely low, providing sufficient contrast. Abnormalities were depicted early on the $^{111}\text{In-leukocyte}$ images due to the relatively fast blood clearance. A major advantage of $^{111}\text{In-liposomes}$ is the ease of preparation of a high-quality radiopharmaceutical without the need to isolate and handle blood.

The superiority of $^{111}\text{In-leukocytes}$ over labeled gamma

globulin confirms clinical results with labeled gamma globulin (5,24). The abnormalities depicted by the labeled leukocytes scintigraphy were shown to correlate better with disease activity than those observed on labeled gamma globulin images, indicating relatively low sensitivity and specificity of the latter agent (24,25). In this study, this was exemplified by the lower relative uptake of $^{111}\text{In-IgG}$ in affected colon as compared with $^{111}\text{In-leukocytes}$ (and $^{111}\text{In-liposomes}$). This lower ratio will obviously make it more difficult to differentiate accumulation in inflamed from normal tissue uptake. Moreover, even the limited physiological excretion of labeled gamma globulin in the gut may interfere with adequate evaluation of diseased colon (5).

The mechanism of accumulation in the target area of $^{111}\text{In-leukocytes}$ and the two new radiopharmaceuticals is entirely different (26). Adequate $^{111}\text{In-leukocytes}$ scintigraphy requires viability of the cells with intact chemotactic capacities, since the labeled cells must actively migrate to an inflammatory target similar to unlabeled leukocytes. When damaged, the labeled cells show prolonged margination in the lungs, followed by enhanced clearance by the mononuclear phagocytic system (27). This further emphasizes the need for careful preparation of leukocytes during labeling. Indium-111-liposomes and $^{111}\text{In-IgG}$ both accumulate in inflamed tissue by virtue of increased vascular permeability (26). Thus, prolonged intravascular activity will be beneficial for the degree of uptake in inflammation. This also explains why agents like labeled colloids (also a particulate radiopharmaceutical) do not provide adequate delineation of colitis (28,29). Colloids have a short circulation time and distribute rapidly to liver, spleen and bone marrow without sufficient focal uptake in disease. Similar characteristics can be observed with non-PEGylated larger liposomes (6,7). Whether any specific uptake of labeled liposomes in inflammatory cells plays a role in accumulation of the radiopharmaceutical in inflammatory foci remains to be established.

CONCLUSION

Indium-111-liposomes were superior to $^{111}\text{In-leukocytes}$ for scintigraphic evaluation of acute colitis due to high focal uptake and relatively low accumulation in many organs. In addition, in view of the ease of preparation, this new formulation is particularly attractive for progression to clinical studies. Indium-111n-IgG was not very well suited for imaging colitis, because the uptake in diseased segments was relatively low and




FIGURE 3. CI (mean \pm s.e.m.) of the radiopharmaceuticals as compared with the macroscopic severity of abnormalities in the affected colon. (Grade 0 = normal; Grade 1 = inflammation; Grade 2 = ulcers.)

adequate delineation of the abnormalities was more difficult than with ^{111}In -leukocytes or ^{111}In -liposomes.

ACKNOWLEDGMENTS

We thank Mr. G. Grutters and Mr. H. Eikholt (Central Animal Laboratory, University of Nijmegen, The Netherlands) for their help in performing the study. The study was supported by a grant from the Foundation of Technical Sciences (Stichting voor de Technische Wetenschappen, STW), The Netherlands. The polyethylene glycol 1900 derivative of distearoyl phosphatidylethanolamine was a gift from Liposome Technology, Inc. (Menlo Park, CA). Cholesterol was from Sigma Chemical Co. (St. Louis, MO) and glutathione from E. Merck (Darmstadt, Germany).

REFERENCES

1. Crama-Bohouth GE, Pena AS, Arndt JW, et al. Value of indium-111 tropolonate autologous granulocyte scintigraphy in the assessment of inflammatory bowel disease. *Scand J Gastroenterol* 1990;178(suppl):93-98.
2. Lange JMA, Boucher CAB, Hollak CEM, et al. Failure of zidovudine prophylaxis after accidental exposure to HIV-1. *N Engl J Med* 1990;322:1375-1377.
3. Rojas-Burk J. Health officials reactions to infection mishaps. *J Nucl Med* 1992;33:13N-27N.
4. Oyen WJG, Claessens RAMJ, van der Meer JWM, Corstens FHM. Detection of subacute infectious foci with indium-111-labeled autologous leukocytes and with indium-111-labeled human nonspecific immunoglobulin G: a prospective comparative study. *J Nucl Med* 1991;32:1854-1860.
5. Oyen WJG, Claessens RAMJ, van der Meer JWM, Rubin RH, Strauss HW, Corstens FHM. Indium-111-labeled human nonspecific immunoglobulin G: a new radiopharmaceutical for imaging infectious and inflammatory foci. *Clin Infect Dis* 1992;14:1110-1119.
6. Morgan JR, Williams LA, Howard CB. Technetium-labeled liposome imaging for deep-seated infection. *Br J Radiol* 1985;58:35-39.
7. O'Sullivan MM, Powell N, French AP, Williams KE, Morgan JR, Williams BD. Inflammatory joint disease: a comparison of liposome scanning, bone scanning and radiography. *Ann Rheum Dis* 1988;47:485-491.
8. Karlowsky JA, Zhanel GG. Concepts on the use of liposomal antimicrobial agents: applications for aminoglycosides. *Clin Infect Dis* 1992;15:654-667.
9. Boerman OC, Storm G, Oyen WJG, et al. Sterically stabilized liposomes labeled with indium-111 to image focal infection. *J Nucl Med* 1995;35:1639-1644.
10. Bakker-Woudenberg IAJM, Lokerse AF, ten Kate MT, Mouton JW, Woodle MC, Storm G. Liposomes with prolonged blood circulation and selective localization in *Klebsiella pneumoniae*-infected lung tissue. *J Infect Dis* 1993;168:164-171.
11. Storm G, Belliot SO, Daemen T, Lasic DD. Surface modification of nanoparticles to oppose uptake by the mononuclear phagocyte system. *Adv Drug Delivery Res* 1995;17:31-48.
12. Allgayer H, Deschryver K, Stenson WF. Treatment with 16,16'-dimethyl prostaglandin E_2 before and after induction of colitis with trinitrobenzenesulfonic acid in rats decreases inflammation. *Gastroenterology* 1989;96:1290-1300.
13. Percy WH, Buron MB, Rose K, Donovan V, Burakoff R. In vitro changes in the properties of rabbit colonic muscularis mucosae in colitis. *Gastroenterology* 1993;104:369-376.
14. Kim HS, Berstad A. Experimental colitis in animal models. *Scand J Gastroenterol* 1992;27:529-537.
15. Babich JW, Graham W, Barrow SA, et al. Technetium-99m-labeled chemotactic peptides: comparison with indium-111-labeled white blood cells for localizing acute bacterial infection in the rabbit. *J Nucl Med* 1993;34:2176-2181.
16. Oyen WJG, Boerman OC, Storm G, et al. Labeled stealth liposomes in experimental infection: an alternative for leukocyte scintigraphy? *Nucl Med Commun* 1996;17:742-748.
17. Lang J, Vigo-Pelfrey C, Martin F. Liposomes composed of partially hydrogenated egg phosphatidylcholines: fatty acid composition, thermal phase behavior and oxidative stability. *Chem Phys Lipids* 1990;53:91-101.
18. Woodle MC, Mathay KK, Newman MS, et al. Versatility in lipid compositions showing prolonged circulation with sterically stabilized liposomes. *Biochim Biophys Acta* 1992;1105:193-200.
19. Storm G, van Bloois L, Brouwer M, Crommelin DJA. The interaction of cytostatic drugs with adsorbents in aqueous media. The potential implication for liposome preparation. *Biochim Biophys Acta* 1985;818:343-351.
20. Gabizon A, Huberty, Straubinger RM, Price DC, Papahadjopoulos D. An improved method for in vivo tracing and imaging of liposomes using a gallium 67-deferoxamine complex. *J Liposome Res* 1988;1:123-135.
21. Hnatowich DJ, Childs RL, Lanteigne D, Najafi A. The preparation of DTPA-coupled antibodies radiolabeled with metallic radionuclides: an improved method. *J Immunol Methods* 1983;65:147-157.
22. Lantto E, Jarvi K, Krekela I, Lantto T, Taavitsainen M, Vedenkangas H, Vorne M. Technetium-99m-hexamethylpropylene amine oxine leukocytes in the assessment of disease activity in inflammatory bowel disease. *Eur J Nucl Med* 1992;19:14-18.
23. Thakur ML, McAfee JG. The significance of chromosomal aberrations in indium-111-labeled lymphocytes. *J Nucl Med* 1984;25:922-927.
24. Arndt JW, van der Sluys-Veer A, Blok D, et al. A prospective comparison of technetium-99m-labeled polyclonal human immunoglobulin and indium-111 granulocytes for localization of inflammatory bowel disease. *Acta Radiol* 1992;33:140-144.
25. Oyen WJG, Naber AHJ, Claessens RAMJ, van der Meer JWM, Corstens FHM. Evaluation of inflammatory bowel disease activity with indium-111-labeled human nonspecific immunoglobulin G [Abstract]. *J Nucl Med* 1992;33:919.
26. Oyen WJG, Boerman OC, van der Laken CJ, Claessens RAMJ, van der Meer JWM, Corstens FHM. The uptake mechanisms of inflammation and infection localizing agents. *Eur J Nucl Med* 1996;23:459-465.
27. Saverymuttu SH, Peters AM, Danpure HJ, Reavy HJ, Osman S, Lavender JP. Lung transit of indium-111-labeled granulocytes: relationship to labeling techniques. *Scand J Haematol* 1983;30:151-160.
28. Wheeler JG, Slack NF, Duncan A, Palmer M, Harvey RF. Technetium-99m-nanocolloid imaging in inflammatory bowel disease. *Nucl Med Commun* 1990;11:127-133.
29. Arndt JW, van der Sluys-Veer A, Blok D, et al. Technetium-99m-nanocolloid for the scintigraphic assessment of inflammatory bowel disease in the colon: its value in comparison with indium-111-labeled granulocytes. *Eur J Radiol* 1991;12:30-34.

Improved Wrist Fracture Localization with Digital Overlay of Bone Scintigrams and Radiographs

W. Roolker, M.M.C. Tiel-van Buul, A.H. Broekhuizen, A.K. Eikelenboom and E.A. van Royen

Departments of Nuclear Medicine, Surgery and Radiology, Academic Medical Center, Amsterdam, The Netherlands

The diagnosis of scaphoid fracture is often difficult and of interest in traumatology. Because of the low sensitivity of repeated scaphoid radiographs, a bone scintigram is advocated and considered the gold standard. In this study, we tried to simplify the interpretation of the bone scintigram of hand and wrist in localizing the hot spot by the digital overlay of the radiograph and the bone scintigram, using a simple device, in patients after wrist trauma. **Methods:** Twenty-one consecutive patients (22 wrists) with clinically-suspected scaphoid fracture and negative initial radiographs were included. The PA view of the wrist was obtained with the hand of the patient placed in an acrylic device with three lead markers. For the bone scan, a similar device was used with ^{57}Co markers at the same

positions. We called this device the "hand-fix." The PA radiograph was digitized with a videocamera and overlaid on the bone scan. Each bone scan was interpreted twice by each of three observers, one nuclear physician and two residents in nuclear medicine. The first interpretation was made without the digital overlay, and the second was made with the digital overlay. **Results:** The bone scintigrams were positive in the scaphoid, distal radius and in other carpal bones. Out of the 22 bone scans, Observer 1 judged 19 correctly, Observer 2 judged 16 correctly and Observer 3 judged 10 correctly without the digital overlay images. All three observers gave a correct localization in the 22 wrists using the digital overlay images. **Conclusion:** The digital overlay of a radiograph and a bone scintigram, using the hand-fix, simplifies and improves interpreting and localizing the hot spot on bone scintigrams in patients with wrist injuries.

Key Words: wrist; bone scintigraphy; multimodality

J Nucl Med 1997; 38:1600-1603

Received Aug. 1, 1996; revision accepted Nov. 6, 1996.

For correspondence or reprints contact: W. Roolker, MD, Department of Nuclear Medicine, F2Noord, Academic Medical Center, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands.