

Erratum

Due to a production error, Tables 1 and 3 in the article "Technetium-99m-ECD Brain SPECT in Misery Perfusion" by Shimosegawa et al. (*JNM* 1997;38:791-796) were printed incorrectly. The tables are reprinted correctly here.

TABLE 1
Summary of Patient Data

| Patient no. | Age (yr) | Sex | Angiographical finding | Main symptom | Interval from initial attack to SPECT-PET study (days) |
|-------------|----------------|-----|------------------------|------------------|--|
| 1 | 69 | M | rt. MCA occlusion | Hemiparesis | 48 |
| 2 | 67 | M | rt. MCA occlusion | Dysarthria | 59 |
| 3 | 72 | M | rt. ICA stenosis | Hemiparesis | 65 |
| 4 | 59 | M | rt. ICA occlusion | Hemiparesis | 140 |
| 5 | 47 | F | rt. MCA occlusion | Dysarthria | 19 |
| 6 | 43 | M | rt. MCA stenosis | Hemiparesis | 11 |
| 7 | 76 | F | lt. MCA stenosis | Dysarthria | 101 |
| 8 | 77 | F | rt. MCA occlusion* | Hemiparesis | 22 |
| 9 | 68 | F | lt. MCA occlusion | Numbness | 49 |
| 10 | 26 | M | rt. ICA occlusion | Hemiparesis | 5 |
| 11 | 36 | M | rt. ICA occlusion | Dysarthria | 12 |
| 12 | 80 | M | rt. ICA occlusion* | Gait disturbance | 12 |
| Average | 60.0 ± 17.0 yr | | | | 45.3 ± 39.7 days |

*Angiographical findings were obtained by magnetic resonance angiography.

TABLE 3
Relationship between Relative Technetium-99m-ECD Activity and Relative PET Values

| SPECT scan time (min) | Patient no. | rCBF | | rCMRO ₂ | |
|-----------------------|-------------|-------|--------------|--------------------|--------------|
| | | r | Significance | r | Significance |
| 1 | 24 | 0.674 | p < 0.001 | 0.569 | p < 0.01 |
| 7 | 24 | 0.561 | p < 0.01 | 0.569 | p < 0.01 |
| 15 | 22 | 0.493 | p < 0.05 | 0.758 | p < 0.001 |
| 25 | 20 | 0.574 | p < 0.01 | 0.815 | p < 0.001 |
| 35 | 22 | 0.567 | p < 0.01 | 0.814 | p < 0.001 |
| 45 | 24 | 0.521 | p < 0.01 | 0.761 | p < 0.001 |
| 55 | 18 | 0.531 | p < 0.05 | 0.789 | p < 0.001 |

r = correlation coefficient.