

## Effect of Obesity on Red Cell Mass

**TO THE EDITOR:** Leslie et al. (1) are certainly correct in emphasizing that obesity will cause errors in calculating predicted blood volumes if normal values of red cell mass are calculated as a linear function of body weight alone. However, in their conclusion that "Commonly used reference ranges generate inconsistent results when extrapolated to obese patients," the authors overlook the most recent recommendation for calculating normal blood volume, which takes obesity into account and offers a more consistent reference range. The authors refer to the 1980 recommendations of the International Committee for Standardization in Haematology (ICSH) (2), which calculate normal blood volumes in terms of body weight. However, they fail to mention the superseding ICSH recommendations of 1995 (3), which are based on a review of several investigations of normal blood volume and consolidation of data that include some obese subjects. That latest report recommends the use of regression equations based on body surface area (calculated from height and weight) to arrive at the predicted normal values for an individual.

### REFERENCES

1. Leslie WD, Dupont JO, Peterdy AE. Effect of obesity on red cell mass results. *J Nucl Med.* 1999;40:422-428.
2. International Committee for Standardization in Haematology. Recommended methods for measurement of red-cell and plasma volume. *J Nucl Med.* 1980;21:793-800.
3. Pearson TC, Guthrie DL, Simpson J, et al. Interpretation of measured red cell mass and plasma volume in adults: Expert Panel on Radionuclides of the International Council for Standardization in Haematology. *Br J Haematol.* 1995;89:748-756.

**Peter B. Schneider**

*U Mass Memorial Medical Center  
Worcester, Massachusetts*

**REPLY:** We thank Dr. Schneider for drawing attention to this omission in our article (1). The updated recommendations of the ICSH (2) are based on a careful reanalysis of pooled data from many of the studies that we considered individually. As such, we agree that they provide the "best" currently available normative regression equations for blood volumes.

### REFERENCES

1. Leslie WD, Dupont JO, Peterdy AE. Effect of obesity on red cell mass results. *J Nucl Med.* 1999;40:422-428.
2. Pearson TC, Guthrie DL, Simpson J, et al. Interpretation of measured red cell mass and plasma volume in adults: Expert Panel on Radionuclides of the International Council for Standardization in Haematology. *Br J Haematol.* 1995;89:748-756.

**William D. Leslie  
Jacqueline O. Dupont  
Anne E. Peterdy**

*St. Boniface General Hospital  
Winnipeg, Manitoba, Canada*

## Mammography and <sup>99m</sup>Tc-MIBI Scintimammography in Suspected Breast Cancer

**TO THE EDITOR:** We read with great interest the article by Prats et al. (1) on the combination of mammography and <sup>99m</sup>Tc-methoxyisobutyl isonitrile (MIBI) scintimammography to prevent unnecessary biopsies in patients with suspected breast cancer. The authors evaluated 90 patients with suspected breast cancer involving 97 lesions and suggested that biopsies should be performed on lesions that have a high suspicion of malignancy as well as those with low or indeterminate suspicion that are smaller than 1 cm or with positive scintimammography results. This protocol would reduce the total number of biopsies by 34%.

We have achieved similar results in our study (2) using a "mamma malignancy index" (MMI) that takes into account the results of scintimammography, mammography, and mammosonography. The results of these modalities were scored from 0 to 2 (0 = negative or low suspicion of malignancy, 1 = indeterminate, and 2 = positive or high probability of malignancy). An MMI was calculated by simply adding the respective scores of each investigation. The prospective study comprised 64 patients with suspect lesions in the breast. Prone breast scintigraphy was performed using the technique of Khalkhali et al. (3). The results suggested a high negative predictive value (100%) for patients with an MMI of <2.

In women with dense breasts, mammography has a false-negative rate of 25%-45% (4), but the combination of mammography and mammosonography is known to increase the number of occult breast cancer cases detected (5). Therefore, we included the results of sonography for calculation of the MMI to characterize a breast lesion as benign and thus prevent unnecessary biopsies. To date, we have examined 94 patients. Biopsies revealed malignancies in 30 patients and benign lesions in 64 patients. Forty-nine patients with an MMI of 0-1 had negative biopsies. Twenty-nine patients had an MMI of 2-3; malignant lesions were revealed in 14 of these patients after biopsy. All patients with an MMI of 4-6 (16/94) had malignancies.

Tumor sizes ranged from 6 to 35 mm in diameter. The negative predictive value in the group of patients with an MMI of 0-1 was again 100%; use of this index could have prevented 52% of the biopsies in this patient sample. The combination of 3 independent imaging modalities established the high indicative value of the MMI. For instance, indeterminate mammography and sonography in 2 patients and indeterminate scintigraphy and sonography in 1 patient indicated correctly the need for biopsy in these 3 patients with malignant breast lesions. On the other hand, negative sonographic results facilitate decision making, which could prevent open biopsies in the benign cases.

In conclusion, we have found that inclusion of mammosonographic results in the MMI improves decisively the discriminatory value of this index by preventing unnecessary biopsies, thus reducing costs and patient inconveniences.

### REFERENCES

1. Prats E, Aisa F, Abós MD, et al. Mammography and <sup>99m</sup>Tc-MIBI scintimammography in suspected breast cancer. *J Nucl Med.* 1999;40:296-301.