
^{123}I -5-IA-85380 SPECT Imaging of Nicotinic Acetylcholine Receptor Availability in Nonsmokers: Effects of Sex and Menstrual Phase

Kelly P. Cosgrove^{1,2}, Effie M. Mitsis^{1,2}, Frederic Bois^{1,2}, Erin Frohlich^{1,2}, Gilles D. Tamagnan^{1,3}, Erica Krantzler^{1,2}, Edward Perry^{1,2}, Paul K. Maciejewski¹, C. Neill Epperson¹, Sharon Allen⁴, Stephanie O'Malley¹, Carolyn M. Mazure¹, John P. Seibyl³, Christopher H. van Dyck¹, and Julie K. Staley^{1,2}

¹Yale University School of Medicine, New Haven, Connecticut; ²Veterans Affairs Connecticut Healthcare System, West Haven, Connecticut; ³Institute for Neurodegenerative Disorders, New Haven, Connecticut; and ⁴University of Minnesota, Minneapolis, Minnesota

The study of the effects of sex and hormones on brain chemistry and neurotransmission is of increasing importance as evidence emerges of sex differences in behavioral symptoms and treatment response in neuropsychiatric disorders. The nicotinic acetylcholine receptor (nAChR) system has been implicated in a variety of psychiatric disorders, including tobacco smoking, for which there is strong evidence supporting sex differences in behaviors and response to smoking cessation treatments. We examined the availability of nAChR containing the β_2 subunit in healthy men and women and the influence of menstrual phase among women. **Methods:** Ten men and 19 women nonsmokers underwent one ^{123}I -5-IA-85380 (^{123}I -5-IA) SPECT scan and one MRI scan. A subset of 9 women, aged 18–39 y, underwent a second ^{123}I -5-IA scan. These 9 women were scanned during the early follicular (days 4–7 in 8 subjects and day 11 in 1 subject) and mid-luteal (days 19–25) phases of their menstrual cycle. Hormone levels were measured in all women to confirm the phase of the cycle. **Results:** Regional brain activity (kBq/cm^3) was higher (39%–54%) in women than in men nonsmokers. When regional brain activity was normalized to total plasma parent to correct for individual differences in radiotracer metabolism (V_T'), differences of 10%–16% were observed, with women greater than men. In contrast, when regional brain activity was normalized to free plasma parent (V_T), there was less than a 4% difference by sex in regional brain β_2 -nAChR availability. These sex differences in kBq/cm^3 and V_T' resulted from significantly higher levels of total plasma parent, free fraction (f_1), and free plasma parent in women than in men nonsmokers. No differences in plasma measures or brain β_2 -nAChR availability were observed across the menstrual cycle for any outcome measure. **Conclusion:** Overall, these findings demonstrate no significant difference in brain β_2 -nAChR availability between men and women nonsmokers or across the menstrual cycle. Importantly, these findings demonstrate sex differences in radiotracer metabolism and plasma protein binding and highlight the critical need to measure plasma radiotracer levels and f_1 in studies that include both sexes.

Key Words: nicotinic; ^{123}I -5-IA-85380; free fraction; sex; menstrual cycle; SPECT

J Nucl Med 2007; 48:1633–1640
DOI: 10.2967/jnumed.107.042317

The importance of studying sex differences in brain neurochemistry has become increasingly evident with the recent demonstration of differences between men and women in symptomatology and treatment for many neuropsychiatric disorders. Evidence of differences in brain structure, function, neurotransmission, and receptor availability between men and women is mounting (1). Sex differences exist in a variety of behaviors related to tobacco smoking, such as craving, sensitivity to nicotine, and response to nicotine replacement therapy (2,3); however, the mechanisms underlying these differences are unclear. Women often experience more difficulty in quitting smoking than men (4–9), and rates of success may differ by menstrual phase. For example, women who quit smoking during the luteal phase of the menstrual cycle reported significantly more tobacco withdrawal and depressive symptoms than did women who quit smoking during the follicular phase, and this effect was related to the attempt to quit, not to a worsening of mood due to phase of cycle (10). These behavioral differences may be due to underlying sex differences in brain chemistry. Several preclinical studies suggest sex differences at the receptor level. The nicotinic acetylcholine receptor (nAChR) containing the β_2 subunit (β_2 -nAChR), one of the initial sites of action of nicotine in the brain and the most critical site for the reinforcing effects of nicotine, is a likely neurochemical substrate mediating these sex differences. Preclinical studies in nicotine-naïve rats (11) and mice (12) suggested higher nAChR numbers in female than in male rats. It is not known whether nicotinic receptor availability differs between living men and women.

Received Mar. 30, 2007; revision accepted Jul. 2, 2007.
For correspondence or reprints contact: Kelly P. Cosgrove, PhD, Yale University School of Medicine and the VACHS, Department of Psychiatry/116A6, 950 Campbell Ave., West Haven, CT 06516.
E-mail: kelly.cosgrove@yale.edu
COPYRIGHT © 2007 by the Society of Nuclear Medicine, Inc.

Menstrual cycle–related fluctuations in the neuroendocrine milieu may contribute to the observed sex differences in smoking-related behaviors. At the receptor level, the actions of sex steroids on nAChRs are complex, and preclinical evidence indicates that progesterone and estrogen have opposing effects on nAChR. Progesterone (13–15) and its A-ring metabolites (14) noncompetitively inhibit nAChR. Progesterone (16) and the neurosteroid pregnenolone (17) also have been shown to block nicotinic receptor response. Conversely, estradiol has been shown to potentiate acetylcholine-evoked currents in human neuronal nAChRs by interacting at the α_4 subunit (18). However, the steroid hormone β -estradiol and others also inhibit acetylcholine-induced currents in rat neurons (19). Thus, there appears to be a relationship between steroid hormones and nAChR numbers. Ovariectomy also reduces coupled acetylcholine synthesis and high-affinity choline uptake, which is reversed by the administration of estradiol in rat brain (20). The effect of estradiol on the state of neuronal nAChRs has also been studied. Chronic estradiol treatment (i.e., 2 mo) in female rats decreases the affinity and increases the binding of ^3H -methylcarbonylcholine, a nicotinic agonist, in the hippocampus and hypothalamus, suggesting that estradiol decreases presynaptic cholinergic function (21).

The development and validation of ^{123}I -5-IA-85380 (^{123}I -5-IA) for brain SPECT allows the *in vivo* examination of β_2 -nAChRs in living men and women. High test–retest reliability and validity were recently reported for ^{123}I -5-IA SPECT (22), and ^{123}I -5-IA SPECT has demonstrated significantly higher β_2 -nAChR availability in recently abstinent tobacco smokers than in age- and sex-matched nonsmokers (23). In the present study, we hypothesized that the sex and cyclical differences in response to nicotine are due to variations in nAChR availability in the brain. This question was addressed by imaging healthy nonsmoking men and women with ^{123}I -5-IA SPECT to examine the effects of sex and menstrual cycle in women on the availability of β_2 -nAChRs.

MATERIALS AND METHODS

Subjects

Study 1: The Effect of Sex on ^{123}I -5-IA Uptake. Ten men (mean age \pm SD, 27.7 ± 7.3 y; range, 20–41 y; 10 Caucasian) and 19 women (mean age, 26.2 ± 6.8 y; range, 20–39 y; 9 Caucasian, 5 Hispanic, 4 African American, and 1 Asian) underwent one ^{123}I -5-IA scan and one MRI scan. Women in study 1 who did not participate in study 2 were scanned during the follicular ($n = 6$) or mid-luteal ($n = 3$) phase of the menstrual cycle, and 1 woman who was taking hormonal contraceptives at the time of the scan was included. Hormones were analyzed to confirm the phase of the cycle and the premenopausal status.

Study 2: The Effect of Menstrual Cycle on ^{123}I -5-IA Uptake. Nine women (mean age, 27.0 ± 7.8 y; range, 18–39 y; 4 white, 2 Hispanic, 2 African American, and 1 Asian) who participated in study 1 were recruited to participate in study 2, which involved undergoing a second ^{123}I -5-IA scan. With the first day of

menstrual flow anchored as day 1, SPECT scans were performed during the early follicular (days 4–7 in 8 subjects and day 11 in 1 subject) and mid-luteal (days 19–25) phases of the menstrual cycle. The subjects were not counterbalanced for the order of follicular versus mid-luteal scans. Of the 9 subjects, 8 completed the 2 scans within 2 consecutive cycles, and 1 completed the scans within 3 consecutive cycles. Hormone levels were measured on scan days to confirm the phase of the cycle. We used estradiol levels of less than 55 pg/mL during the early follicular phase and progesterone levels of more than 3 ng/mL during the mid-luteal phase to confirm cycle phase.

This study was approved by the Yale University School of Medicine Human Investigation Committee, the West Haven Veterans Administration Human Subjects Subcommittee, the Radiation Safety Committee, and the Food and Drug Administration. Subjects were recruited from the community by word of mouth, posters, or newspaper advertisements. Eligibility was determined as follows. All subjects received a medical examination by a study physician to exclude any major medical issues or neurologic disorders. This medical examination included a physical examination, electrocardiogram, serum chemistries, thyroid function studies, complete blood count, urinalysis, and urine toxicology screening. The subjects were interviewed using the Structured Clinical Interview of the *Diagnostic and Statistical Manual of Mental Disorders* to rule out any Axis I disorder. All subjects were never smokers (defined as <100 cigarettes in a lifetime) and had no history of significant medical illness or major head trauma. Nonsmoking status was confirmed by plasma cotinine levels of less than 15 ng/mL, urine cotinine levels of less than 100 ng/mL and carbon monoxide levels of less than 11 ppm on the day of intake and the day of the scan. All women of childbearing age were required to have a negative pregnancy test during the screening process and before radiotracer injection on each study day. All women participating in study 2 were required to have no gynecologic problems and have a history of regular menstrual cycles 26–32 d in length. In the menstrual cycle study, use of hormonal contraceptives within the previous 6 mo was exclusionary.

MRI

MRI was performed on a 1.5-T camera (Sonata; Siemens) in a standard orientation (5- to 7-ms echo time, 24-ms repetition time, 256×192 matrix, 1 excitation, 30-cm field of view, and 124 contiguous slices with a 1.2-mm thickness), and the MRI images were used for coregistration to the SPECT images to provide an anatomic guide for placement of the regions of interest.

^{123}I -5-IA SPECT Scan

All subjects received a 0.6-g saturated solution of potassium iodide in the hour before radiotracer administration. ^{123}I -5-IA was synthesized as previously described (24) and administered as a bolus to constant infusion at a ratio of 7.0 for 8 h. In study 1, men were injected with a bolus of 150.6 ± 23.6 MBq and a continuous infusion of 21.6 ± 3.4 MBq/h and women with a bolus of 152.6 ± 18.5 MBq and a continuous infusion of 22.1 ± 2.9 MBq/h. Thus, the total dose was 345.1 ± 54.3 MBq for men and 349.6 ± 41.2 MBq for women. In study 2, women were injected in the follicular phase with a bolus of 146.8 ± 26.2 MBq and a continuous infusion of 22.4 ± 2.7 MBq/h and in the mid-luteal phase with a bolus of 160.7 ± 8.1 MBq and a continuous infusion of 23.3 ± 0.06 MBq/h. Thus, the total dose was 333.5 ± 56.6 MBq in the follicular phase and 363.8 ± 15.6 MBq in the mid-luteal phase.

Three consecutive 30-min emission scans and one 15-min simultaneous transmission and emission scan were obtained between hours 6 and 8 on a PRISM 3000 XP (Picker) SPECT camera. The PRISM 3000 XP is a 3-head camera equipped with a low-energy, ultra-high-resolution fanbeam collimator (photopeak window, $159 \text{ keV} \pm 10\%$; matrix, 128×128) with a uniform sensitivity across the field of view. A ^{57}Co -distributed source was measured with each experiment to control for day-to-day variation in camera sensitivity. The axial resolution (full width at half maximum) was 12.2 mm, measured with a ^{123}I line source in water in a cylindrical phantom. Blood was drawn before injection and at the beginning and end of the emission scans for analysis of plasma total parent and free fraction (f_1) of parent tracer in plasma.

Image Analysis and Outcome Measures

Images were reconstructed and analyzed as previously described, including nonuniform attenuation correction (22), with a single exception. Specifically, the second SPECT scan for the subjects in study 2 was coregistered to the same position as the first scan in order to apply the same region-of-interest template for each subject. MR images were coregistered to the SPECT images to provide an anatomic guide for placement of the regions of interest using MEDx software (Medical Numerics, Inc.). The chosen regions of interest were those known to contain β_2 -nAChRs and included the frontal, parietal, anterior cingulate, temporal, and occipital cortices; the thalamus, the striatum (an average of caudate and putamen), and the cerebellum. Two raters conducted the analysis. Variability between the raters was less than 12% across regions of interest. The mean of the analysis from the 2 raters is reported.

^{123}I -5IA regional activity (kBq/cm^3) is reported along with the outcome measures V_T' (regional activity divided by total plasma parent between 6 and 8 h), which has shown the best test-retest reliability for ^{123}I -5-IA SPECT (25), and V_T (regional activity divided by free plasma parent between 6 and 8 h), to correct for possible differences in plasma protein binding. Both V_T' and V_T are proportional to the binding potential (binding potential, in mL/g , equals receptor number divided by affinity), which is proportional to the receptor number at equilibrium, given the assumptions that there is no change in affinity and that nondisplaceable (nonspecific and free) uptake does not differ between subjects or between comparison groups. As described previously (25), there is no appropriate reference region for this radiotracer, so nondisplaceable ^{123}I -5-IA uptake could not be measured. Also, to control for differences in metabolism and protein binding, the measures of total plasma parent, f_1 , and free parent, which is defined as total parent $\times f_1$, were compared between groups.

Statistical Analysis

For study 1, preliminary analyses of kBq/cm^3 , V_T' , and V_T revealed that values for these measures were highly correlated between brain regions. Pairwise correlations between brain regions ranged from 0.67 to 0.95 for V_T' , from 0.77 to 0.96 for V_T , and from 0.89 to 0.99 for kBq/cm^3 . Therefore, to obviate issues of multicollinearity in the interpretation of these data, measures of kBq/cm^3 , V_T' , and V_T across brain regions were each reduced to single, main components using principal-component analysis. A contemporary, technical account of principal-component analysis has previously been reported (26). Sex differences in these principal components representing more global measures of

kBq/cm^3 , V_T' , and V_T across brain regions were then examined using unpaired t tests. Bonferroni correction for multiple comparisons across 8 brain regions indicated that P values of less than 0.00625 were highly significant. Because of the high correlations between brain regions, the P values in Table 1 cannot be treated as independent. For study 2, differences in V_T' and V_T between the early follicular and mid-luteal phases were examined with paired t tests.

RESULTS

Study 1: The Effect of Sex

Based on principal-component analyses, single components explained 96% of the variance in kBq/cm^3 , 88% of the variance in V_T' , and 91% of the variance in V_T across brain regions. Sex differences in ^{123}I -5-IA uptake varied by outcome measure (Table 1; Figs. 1 and 2). Across brain regions, the main kBq/cm^3 and V_T' components were significantly greater in women than in men (for kBq/cm^3 : $t_{27} = 4.02$, $P = 0.004$; for V_T' : $t_{27} = 2.20$, $P = 0.0362$). The main V_T component did not differ significantly between women and men.

Statistical analyses (t test) also determined that women have significantly greater f_1 ($t_{27} = 2.50$, $P = 0.02$), total parent ($t_{27} = 2.03$, $P = 0.05$), and plasma free parent ($t_{27} = 2.05$, $P = 0.05$) than do men (Fig. 3).

Study 2: The Effect of Menstrual Cycle

Hormone levels obtained on each SPECT scan day during the early follicular (mean estrogen, 38.1 pg/mL ; mean progesterone, 0.7 ng/mL) and mid-luteal (mean estrogen, 121.1 pg/mL ; mean progesterone, 6.4 ng/mL) phases confirmed the accuracy of the self-reported phase of the cycle. However, although all subjects evidenced a luteal-phase rise in progesterone, 4 subjects did not have luteal-phase progesterone levels greater than 3 ng/mL , suggesting an anovulatory cycle as occurs in as many as 38% of cycles in young women (27) and is consistent with other studies (28). Results did not differ when these subjects were eliminated from the analyses. Radiotracer metabolism and plasma protein binding—for example, total parent and f_1 —did not change between the early follicular and mid-luteal phases of the menstrual cycle (Fig. 4). No significant differences in V_T' were observed in women between the early follicular and mid-luteal phases in any region of interest (thalamus: $t = 0.64$, $P = 0.54$; cortex: $t = 0.48$, $P = 0.64$; striatum: $t = 0.93$, $P = 0.38$, and cerebellum: $t = 1.62$, $P = 0.14$ [data not shown]). Similarly, no significant differences in V_T were observed (thalamus: $t = 0.35$, $P = 0.74$; cortex: $t = 0.26$, $P = 0.80$; striatum: $t = 0.39$, $P = 0.71$ [data for striatum not shown in Fig. 4]; and cerebellum: $t = 0.66$, $P = 0.53$) [Fig. 4]). Additionally, there were no correlations between estrogen or progesterone obtained on scan days and β_2 -nAChR availability using the outcome measure V_T' or V_T .

TABLE 1
Sex Differences Between Men and Women in ¹²³I-5-IA Activity by Outcome Measure

Outcome measure	Men		Women		% Difference*	P
	Mean	SD	Mean	SD		
Regional activity (kBq/cm ³)						
Thalamus	11.6	2.4	17.1	4.7	47	0.0009 [†]
Striatum	5.8	1.7	8.9	2.5	54	0.0007 [†]
Parietal cortex	4.2	0.8	6.3	1.8	49	0.0010 [†]
Frontal cortex	4.6	0.9	6.7	2.0	47	0.0015 [†]
Anterior cingulate	4.6	0.9	6.7	1.9	44	0.0019 [†]
Temporal cortex	5.1	1.0	7.3	2.0	41	0.0023 [†]
Occipital cortex	4.6	0.9	6.5	1.9	41	0.0026 [†]
Cerebellum	5.4	1.1	7.6	2.3	39	0.0054 [†]
V _T '						
Thalamus	46.8	6.6	54.3	9.1	16	0.0151
Striatum	24.5	2.8	27.9	4.0	14	0.0108
Parietal cortex	17.2	2.0	19.9	3.2	16	0.0106
Frontal cortex	18.5	2.6	21.1	3.1	15	0.0145
Anterior cingulate	18.5	2.4	21.0	3.2	13	0.0223
Temporal cortex	20.7	2.7	22.9	3.6	11	0.0489
Occipital cortex	18.6	2.2	20.6	3.3	11	0.0457
Cerebellum	21.6	2.8	23.8	4.1	10	0.0757
V _T						
Thalamus	152.4	29.4	154.7	26.3	2	0.4130
Striatum	79.5	14.0	79.8	12.7	0	0.4787
Parietal cortex	56.0	10.5	57.4	11.8	2	0.3824
Frontal cortex	60.2	12.0	60.8	11.2	1	0.4445
Anterior cingulate	60.0	10.1	60.2	10.2	0	0.4743
Temporal cortex	67.6	11.9	65.6	11.6	-3	0.3372
Occipital cortex	60.6	10.7	59.0	9.9	-3	0.3491
Cerebellum	70.8	13.4	68.0	11.6	-4	0.2770

* $(\text{Women} - \text{men})/\text{men} \times 100$.

[†]P values < 0.00625 are significant after Bonferroni correction.

DISCUSSION

This SPECT study examined the effect of sex and menstrual cycle phase on regional brain uptake of ¹²³I-5-IA. The most notable finding was the statistically significant sex differences in total parent (unmetabolized ¹²³I-5-IA) and in f_1 , suggesting that the radiotracer ¹²³I-5-IA is metabolized differently between men and women and that the amount of ¹²³I-5-IA bound to plasma proteins differs between men and women. Importantly, if not corrected for in the brain outcome measures, these differences may lead to the appearance of a sex difference in nAChR availability. This appearance was clearly demonstrated in the present dataset by the profound sex difference in uncorrected (kBq/cm³) measures of regional brain uptake (39%–54%), as compared with the outcome measure that corrected for metabolism only (but not protein binding), V_T' (10%–16%).

A higher f_1 is associated with a higher radiotracer uptake by the brain (29); thus, because V_T' does not correct for protein binding, the sex difference is due to a higher ¹²³I-5-IA uptake in women than in men but not to a greater β_2 -nAChR availability. Accordingly, there was no difference in β_2 -nAChR availability as evaluated by the outcome measure (V_T) that corrected for both radiotracer metabolism

and the f_1 of radiotracer. Radiotracer metabolism, plasma protein binding, and brain β_2 -nAChR availability did not differ between the early follicular and mid-luteal phases of the menstrual cycle in women nonsmokers.

The Effect of Sex on β_2 -nAChR Availability

This study found no difference in β_2 -nAChR availability between healthy men and women nonsmokers. Preclinical studies indicated that nicotine-naïve female animals had higher global nAChR numbers than did male animals (30,31); however, this finding could not be generalized to humans. Differences between species and between radioligands may account for the discrepancy. Differences in nAChR distribution between species have been reported (32); and the radioligands used, ³H-cytisine and (\pm)-*exo*-2-(2-¹²³I-iodo-5-pyridyl)-7-azabicyclo[2.2.1]heptane), an analog of epibatidine, do not selectively measure the β_2 subtype of nAChR. Additionally, it is likely that radioligand metabolism or protein binding were different between male and female animals. Unfortunately, the small size and blood volume of mice and rats precludes appropriate plasma analysis.




FIGURE 1. $^{123}\text{I-5-IA}$ activity is shown in 10 men (○) and 19 women (●) in 3 selected brain regions: thalamus, cortex (average of cortical regions), and cerebellum. Top row depicts data as kBq/cm^3 uncorrected for metabolism or protein binding, middle row is V_T' , and bottom row is V_T . * P values < 0.00625 are significant after Bonferroni correction.

In an earlier preliminary analysis of 12 subjects (6 men and 6 women), we reported a sex difference in β_2 -nAChR availability (33). These analyses were based on the outcome measure V_T' because the difference in total parent and f_1 was not statistically different and because V_T' had

been shown to be a more reliable and less variable outcome measure than V_T (34). As reported here, when the sample size increased to 29 (19 women and 10 men), the sex difference in f_1 and total parent became significant and a contributing factor to the observed sex difference in brain β_2 -nAChR availability. Specifically, in this study, significant sex differences were found, with higher f_1 and total and free parent in women than in men. The significant difference in f_1 led us to evaluate the data with the outcome measure V_T , which indicated that there is no sex difference in β_2 -nAChR availability between nonsmoking men and women.

A limitation of many neuroreceptor imaging studies is that f_1 often is not measured or reported. It is therefore difficult to determine whether apparent differences in receptor availability between groups may actually be due to a difference in available unbound radiotracer in plasma. Some sex differences in the metabolism of radiotracers do exist. For example, men had a significantly greater percentage of unmetabolized ^{18}F -altanserin in plasma, and thus, women metabolize ^{18}F -altanserin faster than men (35), which led to potentially discrepant findings regarding sex differences in the serotonin 2A receptor (35,36). Specifically, the study that reported sex differences (36) did not




FIGURE 2. Mean parametric images illustrating $^{123}\text{I-5-IA}$ activity in 10 men and 19 women in V_T' and V_T .




FIGURE 3. Radiotracer metabolism and protein binding are shown in 10 men (○) and 19 women (●).

correct for metabolism of ^{18}F -altanserin, and when metabolism was accounted for, no sex difference was found (35). No sex difference in f_1 between men and women was specifically reported for ^{18}F -altanserin (35) and for ^{11}C -WAY-100635 (37). The reason for a sex difference in f_1 in this study is unclear. In humans, both age and sex influence cytochrome p450 enzymatic activity. For example, sex and menstrual phase influence cytochrome p450 3A activity, with women having higher activity than men, specifically in the preovulatory phase (38); however, it has been suggested that the effect of sex is more limited than the effect of intersubject variation (39). There may be a sex difference in protein binding, or it may be that sex-steroid hormones in

women influence protein binding. One study reported that young adult women have lower basal protein oxidation than men (40), but no sex differences were reported for protein turnover (40), plasma protein secretory rates (40), or albumin concentrations (41,42). There was no evidence in the current study that estrogen or progesterone levels were associated with f_1 .

One limitation of this study was that we could not determine the effect of race on β_2 -nAChR availability. There are differences in the metabolism of nicotine both by race (43) and by sex (44,45); thus, it is possible that race may also affect the metabolism and plasma protein binding of ^{123}I -5-IA. Unfortunately, a stringent analysis of race




FIGURE 4. (Top) Within-subject ^{123}I -5-IA activity (V_T) in thalamus, cortex (average of cortical regions), and cerebellum in 9 nonsmoking women in early follicular and mid-luteal phases of menstrual cycle. (Bottom) Radiotracer metabolism and protein binding in same women in early follicular (○) and mid-luteal (●) phases of menstrual cycle. No significant differences in V_T , total parent, f_1 , or free parent were found between early follicular and mid-luteal phases of menstrual cycle.

requires a larger sample and is beyond the scope of this study.

Another limitation of this study was that we examined men and women nonsmokers under the age of 45 y. Thus, we do not know yet if the sex differences in metabolism and protein binding of the radiotracer persist across the life span. We chose this age range to focus on a population of premenopausal women, because we had hypothesized that the sex differences in β_2 -nAChR availability were due to the sex-steroid hormones β -estradiol and progesterone, which vary over the course of the menstrual cycle. Although a finding of no sex difference in β_2 -nAChR availability in nonsmokers does not preclude a sex difference in β_2 -nAChR availability in smokers, it does raise the question of whether availability of β_2 -nAChRs is involved in the sex differences in tobacco-smoking behavior, such as differences in quitting smoking and in response to nicotine replacement therapy. Although ^{123}I -5-IA SPECT measures the availability of β_2 -nAChR, this measure may be confounded by occupancy of the receptor by endogenous acetylcholine (46). However, no reports of sex differences in endogenous acetylcholine levels could be found, and plasma acetylcholine does not vary in women with gonadal hormone levels (47). Also, it is not clear if ^{123}I -5-IA is measuring only cell-surface receptors or also internalized receptors; thus, these findings do not rule out sex differences in receptor internalization or synthesis. Finally, other nAChR subtypes may underlie behavioral sex differences. For example, a study found sex differences in β_3 -nAChR expression in the nasal mucosa of human subjects (48). Future studies are needed to evaluate these alternatives.

The Effect of Menstrual Cycle Phase on β_2 -nAChR Availability

This study also found no difference in β_2 -nAChR availability between the early follicular and mid-luteal phases of the menstrual cycle in healthy women nonsmokers. Additionally, in correlational analyses, we observed no relationship between β_2 -nAChR availability and plasma estrogen or progesterone levels or in their ratio. These results suggest that sex-steroid hormones do not acutely regulate β_2 -nAChRs. Alternatively, because we imaged during the early follicular phase, when estrogen and progesterone levels are low, and the mid-luteal phase, when both estrogen and progesterone levels are high, we cannot rule out the possibility that estrogen and progesterone have opposing regulatory effects on β_2 -nAChR availability. Preclinical data suggest an inhibitory role of progesterone and a possible potentiating effect of estrogen on nAChR; thus, during the mid-luteal phase, the high levels of progesterone may be directly inhibiting nAChR or negating the potentiating effects of estrogen on nAChR. β_2 -nAChR availability may appear similar between the early follicular and mid-luteal phases but may be increased in the absence of progesterone, when only estrogen levels are high—for

example, the late follicular phase. One limitation of having studied women during the early to mid-follicular and mid-luteal phases of the menstrual cycle is that the relative effects of estradiol and progesterone on β_2 -nAChR could not be discerned. The influence of menstrual phase and sex-steroid hormones on other neurotransmitter and receptor systems has been underresearched. Striatal dopamine transporters do not vary with menstrual cycle phase (28), and there is conflicting evidence on variations in dopamine D_2 receptors by phase of cycle (49,50). Serotonin transporter availability also does not fluctuate (28), whereas the serotonin 2A receptor (51) may fluctuate with menstrual cycle phase. Finally, γ -aminobutyric acid levels were tightly regulated by the female menstrual cycle (52). Generally, sex steroids may not have acute effects on receptor numbers in the brain, but effects may be more likely to occur in utero, during other developmental periods such as adolescence, or over time because of chronic exposure to sex-steroid hormones.

CONCLUSION

Few in vivo PET or SPECT neuroreceptor imaging studies have examined sex differences or changes in neurotransmitter levels or receptors across the menstrual cycle in living humans, and there are virtually no data on the effects of menstrual phase on human brain derived from postmortem studies. Studies examining the effect of sex and menstrual phase on the brain are important because many psychiatric disorders are marked by sex differences in signs and symptoms that may be a consequence of sex differences in brain chemistry. Here, we have demonstrated no significant difference in brain β_2 -nAChR availability between men and women nonsmokers or across the menstrual cycle. Importantly, these findings demonstrate sex differences in radiotracer metabolism and plasma protein binding and highlight the critical need to measure plasma radiotracer levels and f_1 in studies that include both sexes.

ACKNOWLEDGMENTS

This work was supported by grants from NIH (RO1DA 015577, P50AA15632, KO1DA020651, KO2DA021863, and NS044316) and by the Ethel F. Donaghue Women's Health Investigator Program of Women's Health Research at Yale. We thank Louis Amici, Nina Sheung, Meredith Terlecki, Tracy Kloczynski, Stephanie Stiklus, and Noah McGuire-Schwartz for technical assistance.

REFERENCES

1. Cosgrove KP, Mazure CM, Staley JK. Evolving knowledge of sex differences in brain structure, function, and chemistry. *Biol Psychiatry*. June 1, 2007 [Epub ahead of print].
2. Perkins KA, Donny E, Caggiula AR. Sex differences in nicotine effects and self-administration: review of human and animal evidence. *Nicotine Tob Res*. 1999;1:301–315.
3. Perkins K. Nicotine discrimination in men and women. *Pharmacol Biochem Behav*. 1999;64:295–299.

4. Bjornson W, Rand C, Connett JE, et al. Gender differences in smoking cessation after 3 years in the Lung Health Study. *Am J Public Health.* 1995;85:223–230.
5. Wetter D, Kenford S, Smith S, Fiore M, Jorenyb D, Baker T. Gender differences in smoking cessation. *J Consult Clin Psychol.* 1999;67:555–562.
6. Perkins K. Sex differences in nicotine vs. non-nicotine reinforcement as determinants of tobacco smoking. *Exp Clin Psychopharmacol.* 1996;4:166–177.
7. Osler M, Prescott E, Godtfredsen N, Hein H, Schnohr P. Gender and determinants of smoking cessation: a longitudinal study. *Prev Med.* 1999;29:57–62.
8. Sherman SE, Fu SS, Joseph AM, Lanto AB, Yano EM. Gender differences in smoking cessation services received among veterans. *Womens Health Issues.* 2005;15:126–133.
9. Killen J, Fortmann S, Varady A, Kraemer H. Do men outperform women in smoking cessation trials? Maybe, but not by much. *Exp Clin Psychopharmacol.* 2002;10:295–301.
10. Perkins KA, Levine M, Marcus M, et al. Tobacco withdrawal in women and menstrual cycle phase. *J Consult Clin Psychol.* 2000;68:176–180.
11. Koylu E, Demireoren S, London E, Pogun S. Sex difference in up-regulation of nicotinic acetylcholine receptors in rat brain. *Life Sci.* 1997;61:PL185–PL190.
12. Mochizuki T, Villemagne V, Scheffel U, et al. Nicotine induced up-regulation of nicotinic receptors in CD-1 mice demonstrated with an in vivo radiotracer: gender differences. *Synapse.* 1998;30:116–118.
13. Valera S, Ballivet M, Bertrand D. Progesterone modulates a neuronal nicotinic acetylcholine receptor. *Proc Natl Acad Sci USA.* 1992;89:9949–9953.
14. Bullock A, Clark A, Grady S, et al. Neurosteroids modulate nicotinic receptor function in mouse striatal and thalamic synaptosomes. *J Neurochem.* 1997;68:2412–2423.
15. Ke L, Lukas R. Effects of steroid exposure on ligand binding and functional activities of diverse nicotinic acetylcholine receptor subtypes. *J Neurochem.* 1996;67:1100–1112.
16. Takashima K, Kawasaki S, Kimura S, Fujita R, Sasaki K. Blockade of ionotropic receptor responses by progesterone in the ganglion cells of *Aplysia*. *Neurosci Res.* 2002;43:119–125.
17. Kudo K, Tachikawa E, Kashimoto T. Inhibition by pregnenolone sulfate of nicotinic acetylcholine response in adrenal chromaffin cells. *Eur J Pharmacol.* 2002;456:19–27.
18. Curtis L, Buisson B, Bertrand S, Bertrand D. Potentiation of human alpha4beta2 neuronal nicotinic acetylcholine receptor by estradiol. *Mol Pharmacol.* 2002;61:127–135.
19. Uki M, Nabekura J, Akaike N. Suppression of the nicotinic acetylcholine response in rat superior cervical ganglionic neurons by steroids. *J Neurochem.* 1999;72:808–814.
20. O'Malley CA, Hautamaki RD, Kelley M, Meyer EM. Effects of ovariectomy and estradiol benzoate on high affinity choline uptake, ACh synthesis, and release from rat cerebral cortical synaptosomes. *Brain Res.* 1987;403:389–392.
21. Lapchak PA, Araujo DM, Quirion R, Beaudet A. Chronic estradiol treatment alters central cholinergic function in the female rat: effect on choline acetyltransferase activity, acetylcholine content, and nicotinic autoreceptor function. *Brain Res.* 1990;525:249–255.
22. Staley JK, van Dyck CH, Weinzimmer D, et al. ¹²³I-5-IA-85380 SPECT measurement of nicotinic acetylcholine receptors in human brain by the constant infusion paradigm: feasibility and reproducibility. *J Nucl Med.* 2005;46:1466–1472.
23. Staley JK, Krishnan-Sarin S, Cosgrove KP, et al. Human tobacco smokers in early abstinence have higher levels of beta2* nicotinic acetylcholine receptors than nonsmokers. *J Neurosci.* 2006;26:8707–8714.
24. Zoghbi S, Tamagnan G, Fujita M, et al. Measurement of plasma metabolites of (S)-5-[¹²³I]iodo-3-(2-azetidylmethoxy) pyridine (5-IA-85380), a nicotinic acetylcholine receptor imaging agent, in non-human primates. *Nucl Med Biol.* 2001;28:91–96.
25. Staley JK, van Dyck C, Weinzimmer D, et al. ¹²³I-5-IA-85380 SPECT measurement of nicotinic acetylcholine receptors in human brain by the constant infusion paradigm: feasibility and reproducibility. *J Nucl Med.* 2005;46:1466–1472.
26. Jolliffe I. *Principal Components Analysis.* New York, NY: Springer-Verlag; 2002.
27. Metcalf MG, Mackenzie JA. Incidence of ovulation in young women. *J Biosoc Sci.* 1980;12:345–352.
28. Best SE, Sarrel PM, Malison RT, et al. Striatal dopamine transporter availability with [¹²³I]beta-CIT SPECT is unrelated to gender or menstrual cycle. *Psychopharmacology (Berl).* 2005;183:181–189.
29. Laruelle M, Slifstein M, Huang Y. Relationships between radiotracer properties and image quality in molecular imaging of the brain with positron emission tomography. *Mol Imaging Biol.* 2003;5:363–375.
30. Mochizuki T, Villemagne VL, Scheffel U, et al. Nicotine induced up-regulation of nicotinic receptors in CD-1 mice demonstrated with an in vivo radiotracer: gender differences. *Synapse.* 1998;30:116–118.
31. Koylu E, Demireoren S, London ED, Pogun S. Sex difference in up-regulation of nicotinic acetylcholine receptors in rat brain. *Life Sci.* 1997;61:PL185–PL190.
32. Gotti C, Clementi F. Neuronal nicotinic receptors: from structure to pathology. *Prog Neurobiol.* 2004;74:363–396.
33. Staley J, Cosgrove K, Epperson N, et al. Imaging sex differences in beta2-nicotinic acetylcholine receptor expression in nonsmokers using [¹²³I]-5-IA SPECT. Paper presented at: 2005 meeting of the Society for Research on Nicotine and Tobacco; March 22, 2005; Prague, Czech Republic.
34. Staley J, van Dyck CH, Weinzimmer D, et al. Iodine-¹²³-5-IA-85380 SPECT measurement of nicotinic acetylcholine receptors in human brain by the constant infusion paradigm: feasibility and reproducibility. *J Nucl Med.* 2005;46:1466–1472.
35. Adams K, Pinborg L, Svarer C, et al. A database of [¹⁸F]-altanserin binding to 5-HT_{2A} receptors in normal volunteers: normative data and relationship to physiological and demographic variables. *Neuroimage.* 2004;21:1105–1113.
36. Biver F, Lotstra F, Monclus M, et al. Sex difference in 5HT₂ receptor in the living human brain. *Neurosci Lett.* 1996;204:25–28.
37. Parsey R, Oquendo M, Simpson N, et al. Effects of sex, age, and aggressive traits in man on brain serotonin 5-HT_{1A} receptor binding potential measured by PET using [C-11]WAY-100635. *Brain Res.* 2002;954:173–182.
38. Zhu B, Liu ZQ, Chen GL, et al. The distribution and gender difference of CYP3A activity in Chinese subjects. *Br J Clin Pharmacol.* 2003;55:264–269.
39. Bebia Z, Buch SC, Wilson JW, et al. Bioequivalence revisited: influence of age and sex on CYP enzymes. *Clin Pharmacol Ther.* 2004;76:618–627.
40. Volpi E, Lucidi P, Bolli GB, Santeusano F, De Feo P. Gender differences in basal protein kinetics in young adults. *J Clin Endocrinol Metab.* 1998;83:4363–4367.
41. Niemela JE, Csako G, Bui MN, Elin RJ. Gender-specific correlation of platelet ionized magnesium and serum low-density-lipoprotein cholesterol concentrations in apparently healthy subjects. *J Lab Clin Med.* 1997;129:89–96.
42. Mattix HJ, Hsu CY, Shaykevich S, Curhan G. Use of the albumin/creatinine ratio to detect microalbuminuria: implications of sex and race. *J Am Soc Nephrol.* 2002;13:1034–1039.
43. Perez-Stable E, Herrera B, Jacob P III, Benowitz N. Nicotine metabolism and intake in black and white smokers. *JAMA.* 1998;280:152–156.
44. Benowitz NL, Lessov-Schlaggar CN, Swan GE, Jacob P III. Female sex and oral contraceptive use accelerate nicotine metabolism. *Clin Pharmacol Ther.* 2006;79:480–488.
45. Hukkanen J, Jacob P III, Benowitz NL. Metabolism and disposition kinetics of nicotine. *Pharmacol Rev.* 2005;57:79–115.
46. Fujita M, Al-Tikriti M, Tamagnan G, et al. Influence of acetylcholine levels on the binding of a SPECT nicotinic acetylcholine receptor ligand [¹²³I]-5-IA-85380. *Synapse.* 2003;48:116–122.
47. Kawashima K, Oohata H, Fujimoto K, Suzuki T. Plasma concentration of acetylcholine in young women. *Neurosci Lett.* 1987;80:339–342.
48. Keiger CJ, Case LD, Kendal-Reed M, Jones KR, Drake AF, Walker JC. Nicotinic cholinergic receptor expression in the human nasal mucosa. *Ann Otol Rhinol Laryngol.* 2003;112:77–84.
49. Nordstrom AL, Olsson H, Halldin CA. PET study of D2 dopamine receptor density at different phases of the menstrual cycle. *Psychiatry Res.* 1998;83:1–6.
50. Wong DF, Broussolle EP, Wand G, et al. In vivo measurement of dopamine receptors in human brain by positron emission tomography: age and sex differences. *Ann N Y Acad Sci.* 1988;515:203–214.
51. Wihlback AC, Sundstrom Poromaa I, Bixo M, Allard P, Mjorndal T, Spigset O. Influence of menstrual cycle on platelet serotonin uptake site and serotonin_{2A} receptor binding. *Psychoneuroendocrinology.* 2004;29:757–766.
52. Epperson CN, Haga K, Mason GF, et al. Cortical gamma-aminobutyric acid levels across the menstrual cycle in healthy women and those with premenstrual dysphoric disorder: a proton magnetic resonance spectroscopy study. *Arch Gen Psychiatry.* 2002;59:851–858.