- Ran C, Zhang Z, Hooker J, Moore A. In vivo photoactivation without "light": use of Cherenkov radiation to overcome the penetration limit of light. *Mol Imaging Biol.* 2012;14:156–162.
- Kotagiri N, Sudlow GP, Akers WJ, Achilefu S. Breaking the depth dependency of phototherapy with Cerenkov radiation and low-radiance-responsive nanophotosensitizers. *Nat Nanotechnol.* 2015;10:370–379.
- Kamkaew A, Cheng L, Goel S, et al. Cerenkov radiation induced photodynamic therapy using chlorin e6-loaded hollow mesoporous silica nanoparticles. ACS Appl Mater Interfaces. 2016;8:26630–26637.
- Kavadiya S, Chadha TS, Liu H, Shah VB, Blankenship RE, Biswas P. Directed assembly of the thylakoid membrane on nanostructured TiO₂ for a photo-electrochemical cell. *Nanoscale*. 2016;8:1868–1872.
- de Carvalho BA, Kavadiya S, Huang S, Niedzwiedzki DM, Biswas P. Highly stable perovskite solar cells fabricated under humid ambient conditions. *IEEE Journal of Photovoltaics*. 2017;7:532–538.
- Kavadiya S, Niedzwiedzki DM, Huang S, Biswas P. Electrospray-assisted fabrication of moisture-resistant and highly stable perovskite solar cells at ambient conditions. *Advanced Energy Materials*. 2017. https://doi.org/10.1002/aenm.201700210. Accessed November 19, 2018.
- Lin L-Y, Nie Y, Kavadiya S, Soundappan T, Biswas P. N-doped reduced graphene oxide promoted nano TiO₂ as a bifunctional adsorbent/photocatalyst for CO₂ photoreduction: effect of N species. *Chem Eng J.* 2017;316:449–460.
- Almquist CB, Biswas P. Role of synthesis method and particle size of nanostructured TiO₂ on its photoactivity. J Catal. 2002;212:145–156.
- Namkhang P. An W-J, Wang W-N, Rane KS, Kongkachuichay P, Biswas P. Low temperature synthesis of N-doped TiO₂ nanocatalysts for photodegradation of methyl orange. *J Nanosci Nanotechnol.* 2013;13:2376–2381.
- Duan D, Liu H, Xu Y, et al. Activating TiO₂ nanoparticles: gallium-68 serves as a high-yield photon emitter for Cerenkov-induced photodynamic therapy. ACS Appl Mater Interfaces. 2018;10:5278–5286.
- Pratt EC, Shaffer TM, Zhang Q, Drain CM, Grimm J. Nanoparticles as multimodal photon transducers of ionizing radiation. *Nat Nanotechnol.* 2018;13:418–426.
- Zhou Z, Song J, Nie L, Chen X. Reactive oxygen species generating systems meeting challenges of photodynamic cancer therapy. *Chem Soc Rev.* 2016;45: 6597–6626.

- Glaser AK, Zhang R, Gladstone DJ, Pogue BW. Optical dosimetry of radiotherapy beams using Cherenkov radiation: the relationship between light emission and dose. *Phys Med Biol.* 2014;59:3789–3811.
- Glaser AK, Zhang R, Andreozzi JM, Gladstone DJ, Pogue BW. Cherenkov radiation fluence estimates in tissue for molecular imaging and therapy applications. *Phys Med Biol.* 2015;60:6701.
- Gill RK, Mitchell GS, Cherry SR. Computed Cerenkov luminescence yields for radionuclides used in biology and medicine. *Phys Med Biol.* 2015;60:4263– 4280.
- Beattie BJ, Thorek DL, Schmidtlein CR, Pentlow KS, Humm JL, Hielscher AH. Quantitative modeling of Cerenkov light production efficiency from medical radionuclides. *PLoS One.* 2012;7:e31402.
- Ouyang Z, Liu B, Yasmin-Karim S, Sajo E, Ngwa W. Nanoparticle-aided external beam radiotherapy leveraging the Čerenkov effect. *Phys Med.* 2016;32:944–947.
- Almquist CB, Biswas P. A mechanistic approach to modeling the effect of dissolved oxygen in photo-oxidation reactions on titanium dioxide in aqueous systems. *Chem Eng Sci.* 2001;56:3421–3430.
- Kotagiri N, Laforest R, Achilefu S. Reply to 'Is Cherenkov luminescence bright enough for photodynamic therapy?'. Nat Nanotechnol. 2018;13:354–355.
- Anderson W, Belcher E. A standard light source of very low intensity based on the Cerenkov effect. *British Journal of Applied Physics*. 1954;5:53–57.
- National Institute of Standards and Technology. ESTAR stopping-power and range tables for electrons. https://physics.nist.gov/PhysRefData/Star/Text/ESTAR. html. Accessed November 19, 2018.
- Physical Sciences Resource Center. Lund/LBNL nuclear data search. https:// psrc.aapt.org/items/detail.cfm?ID=258. Accessed November 19, 2018.
- 36. Villarreal TL, Gómez R, Gonzalez M, Salvador P. A kinetic model for distinguishing between direct and indirect interfacial hole transfer in the heterogeneous photooxidation of dissolved organics on TiO₂ nanoparticle suspensions. *J Phys Chem B*. 2004;108:20278–20290.
- Gerischer H, Heller A. The role of oxygen in photooxidation of organic molecules on semiconductor particles. J Phys Chem. 1991;95:5261–5267.
- Jiang J, Oberdörster G, Elder A, Gelein R, Mercer P, Biswas P. Does nanoparticle activity depend upon size and crystal phase? *Nanotoxicology*. 2008;2:33–42.
- Niu G, Chen X. When radionuclides meet nanoparticles. Nat Nanotechnol. 2018;13:359–360.

Errata

In the article "¹⁸F-Flortaucipir PET/MRI Correlations in Nonamnestic and Amnestic Variants of Alzheimer Disease," by Nasrallah et al. (*J Nucl Med.* 2018;59:299–306), grant numbers were inadvertently left out of the Disclosure section. The Disclosure should have included the following: The study was also partially funded by NIH grant numbers P01 AG017586 (PI. Virginia M. Lee), P30-AG010124 (PI. John Q. Trojanowski), and AG054519 (PI. Murray Grossman). The authors regret the error.

In the article "Immediate Postablation ¹⁸F-FDG Injection and Corresponding SUV Are Surrogate Biomarkers of Local Tumor Progression After Thermal Ablation of Colorectal Carcinoma Liver Metastases," by Cornelis et al. (*J Nucl Med.* 2018;59:1360–1365), values in the SUV_{mean} and SUV_{max} columns of Table 1 were reversed. The corrected table appears below. The authors regret the error.

Average 50VS of hois before and Arter Ablation		
ROI	$\mathrm{SUV}_{\mathrm{mean}}$	$\mathrm{SUV}_{\mathrm{max}}$
Including tumors		
Before ablation	2 (0.6)	9.2 (6.2)
After ablation	2.2 (0.6)	4.5 (1.7)
Including liver background only		
Before ablation	2.5 (0.5)	3.8 (1.6)
After ablation	3.2 (0.6)	4 (0.9)

TABLE 1			
Ave	age SUVs of ROIs Before and	d After Ablation	