

Visualizing T-Cell Responses: The T-Cell PET Imaging Toolbox

Chao Li^{1,2}, Chaozhe Han^{1,2}, Shao Duan^{1,2}, Ping Li³, Israt S. Alam⁴, and Zunyu Xiao^{1,4}

¹Harbin Medical University, Harbin, China; ²Department of Orthopedics, Second Affiliated Hospital of Harbin Medical University, Harbin, China; ³Department of Radiology and Nuclear Medicine, Second Affiliated Hospital of Harbin Medical University, Harbin, China; and ⁴Molecular Imaging Program at Stanford, Department of Radiology, Stanford University School of Medicine, Stanford, California

T lymphocytes are key mediators of the adaptive immune response. Inappropriate or imbalanced T-cell responses are underlying factors in cancer progression, allergy, and other immune disorders. Monitoring the spatiotemporal dynamics of T cells and their functional status has the potential to provide unique biologic insights into health and disease. Noninvasive PET imaging represents an ideal whole-body modality for achieving this goal. With the appropriate PET imaging probes, T-cell dynamics can be monitored in vivo with high specificity and sensitivity. Herein, we provide a comprehensive overview of the applications of this state-of-the-art T-cell PET imaging toolbox and the potential it has to improve the clinical management of cancer immunotherapy and T-cell-driven diseases. We also discuss future directions and prospects for clinical translation.

Key Words: T-cell PET imaging; graft-versus-host disease; inflammatory bowel disease; rheumatoid arthritis; cancer immunotherapy

J Nucl Med 2022; 63:183–188
DOI: 10.2967/jnumed.121.261976

T lymphocytes play a central role in the adaptive immune response. The interplay between T cells and extracellular factors maintains a careful balance between activation, proliferation, survival, and inhibition. Dysregulation of T-cell responses can contribute to cancer progression, allergy, and immune disorders (1,2). Noninvasive PET imaging represents an ideal approach for visualizing T-cell dynamics in vivo, which could improve our understanding of their role in disease pathogenesis. This has motivated the development and evaluation of numerous T-cell imaging probes in both preclinical and clinical settings (Fig. 1). The T-cell PET imaging toolbox consists of a diverse set of approaches that include direct labeling of cells in vitro, proteins and peptides targeting endogenous T-cell surface and secreted biomarkers, small-molecule metabolic tracers, and engineering cells to express PET reporter genes (3). In direct cell labeling, immune cells are incubated with radiolabels ex vivo before adoptive transfer into a living subject for subsequent imaging. Although this is a relatively straightforward methodology for T-cell tracking, its wider clinical

use to date has been limited. Incorporation of radionuclides can cause toxicities, such as radiolysis, and can adversely impact T-cell function. The radiolabel itself becomes diluted as cells divide and proliferate in vivo, reducing the utility of this approach for longitudinal imaging (4). Given the drawbacks of this approach, we will focus our review on the alternative approaches mentioned above, which are summarized in Table 1.

UTILITY OF T-CELL PET IMAGING FOR DIAGNOSIS OF ALLOGENIC AND AUTOIMMUNE DISEASES

Inappropriate T-cell activation and trafficking are seen in a range of pathologies, including acute graft-versus-host disease (GVHD) in the allogeneic transplant setting, and autoimmune diseases such as rheumatoid arthritis (RA) and inflammatory bowel disease. It is well documented that early diagnosis and timely therapeutic intervention in these scenarios can lead to better outcomes and minimize organ damage (5,6), illustrating the urgent need to develop more reliable diagnostic tools. T-cell PET imaging represents an ideal approach for achieving noninvasive, early detection of disease, before the onset of clinical symptoms.

T-cell imaging with 2'-deoxy-2'-¹⁸F-fluoro-9-β-D-arabinofuranosylguanine (¹⁸F-AraG) has successfully detected activated T cells in secondary lymphoid organs at both early and late stages of disease in a murine acute-GVHD model (Fig. 2A) (7) and is now being evaluated clinically in hematopoietic stem cell transplant recipients (NCT03367962). Imaging of the T-cell surface lineage marker CD3 successfully visualized the total T-cell compartment in vivo in a humanized GVHD model (8). However, downregulation of CD3 expression during T-cell activation poses a limitation to this approach. A ⁶⁴Cu-labeled OX40 monoclonal antibody demonstrated excellent diagnostic potential in a murine acute-GVHD model, detecting T-cell activation early in disease and before presentation of overt clinical symptoms (Fig. 2B) (9). However, the agonist OX40 monoclonal antibody clone used in this model also accelerated acute GVHD lethality, even when administered at the relatively low mass doses used for imaging. These findings highlight the importance of testing for biologic perturbations and overall safety before clinical translation. Given the utility of OX40 as a biomarker of GVHD, using an OX40 antagonist clone may be a safer imaging approach for GVHD and other indications for which T-cell activation may be deleterious.

Imaging activated T cells also represents a promising strategy for early diagnosis of RA. Although ¹⁸F-FDG was tested in preclinical and clinical RA studies, the glucose metabolism pathway is shared by multiple tissues, which results in a high rate of

Received Jul. 16, 2021; revision accepted Dec. 6, 2021.
For correspondence or reprints, contact Zunyu Xiao (xiao_zunyu@hrbmu.edu.cn) or Israt S. Alam (israt@stanford.edu).
Published online Dec. 9, 2021.
COPYRIGHT © 2022 by the Society of Nuclear Medicine and Molecular Imaging.




FIGURE 1. Approaches to PET imaging of T cells. T-cell PET imaging toolbox expanded rapidly over last decade. (A) Numerous T-cell-specific PET tracers have been developed, including radiolabeled antibodies and antibody fragments, protein scaffolds, small molecules, and tracers complementary to reporter genes that can be used to track engineered T cells. (B) These approaches have been evaluated in both preclinical and clinical studies. (C) Noninvasive PET imaging of T cells has potential to be highly useful, allowing preclinical researchers and clinicians to predict or monitor therapeutic response to cancer immunotherapy. This modality also enables early diagnosis of inflammatory diseases, such as GVHD, RA, and inflammatory bowel disease, for timely and effective intervention.

false-positives, especially when the region of interest is adjacent to metabolically active tissues (10). Evaluation of ^{18}F -AraG in a pre-clinical adjuvant-induced arthritis model (Fig. 2C) (11) showed significantly higher accumulation of ^{18}F -AraG in RA-affected paws in both the acute and the chronic phases of disease, demonstrating its potential utility for RA diagnosis. Because of the enormous challenge of developing small-molecule binders, antibody-based PET tracers known as immuno-PET probes are favored as a reliable tool for immune cell imaging given their robust and facile radiolabeling protocols. Immuno-PET tracers targeting T-cell surface or secreted biomarkers should also warrant evaluation for RA detection in future studies (12).

For inflammatory bowel disease detection, an anti-CD4 monoclonal antibody labeled with ^{111}In was previously evaluated for SPECT imaging of CD4-positive T cells. More recently, an ^{89}Zr -labeled CD4-specific GK1.5 cys-diabody was developed and evaluated in a mouse model of dextran sulfate sodium-induced colitis (Fig. 2D) (13). Increased accumulation of ^{89}Zr -maleimide-deferoxamine-GK1.5 cDb was detected in the distal colon of colitis mice and further corroborated by increased CD4-positive immunohistochemistry staining, demonstrating the sensitivity of this probe for CD4-positive T cells *in vivo*. Whether PET quantification correlated with disease severity was not discussed in this study; further evaluation is therefore required to explore the feasibility of this approach for determining disease severity. Additionally, naïve CD4-positive T cells residing in the gut may raise the background signal in CD4-targeted imaging. Since

activated T cells are the true mediators of inflammatory bowel disease pathogenesis, PET tracers specific to activated T cells may be preferred.

T-CELL PET IMAGING FOR PREDICTING AND MONITORING IMMUNE RESPONSE TO CANCER IMMUNOTHERAPIES

Immunooncology has rapidly evolved over the last decade (14), with numerous clinical trials demonstrating the unprecedented success of cancer immunotherapies in treating late-stage and recurrent malignancies, including relapsed or refractory B-cell malignancies, melanoma, bladder cancer, and non-small cell lung carcinoma (15–18). Despite the meteoric rise of these innovative therapies, only a small fraction of patients exhibits durable responses, highlighting the urgent need for reliable tools to monitor and predict therapeutic response (19). Anatomic imaging often lacks sensitivity and does not collect direct molecular or functional information about the T-cell compartment. Biopsies are invasive, risk iatrogenic complications, fail to capture whole-body information, and do not assess tumor heterogeneity (20,21). Longitudinal PET imaging and quantitative analysis of T-cell distribution in tumors or secondary lymphoid organs before and after treatment may correlate with therapeutic response, helping clinicians to identify patients most likely to benefit from treatment (22). With this motivation, a wide

range of T-cell-specific PET tracers has been developed and evaluated in preclinical studies and clinical trials.

Endogenous T-cell biomarkers include those specifically expressed on the cell surface, such as T-cell lineage markers (CD4, CD8); those secreted by certain T-cell subsets, such as granzyme B; and activation markers, such as CD278 and CD134 (Figs. 3A and 3B). A major advantage of using immuno-PET probes is their high specificity and affinity toward their homologous target, generating high signal-to-noise ratios and high-contrast images. Neutral binders without agonistic and antagonistic functions have the potential to be widely applicable without concern about perturbative effects. However, because of their large size, intact antibodies (150 kDa) can suffer from poor penetration into target tissues and slow clearance. Additionally, the crystallizable fragment regions of whole antibodies can bind nonspecifically to crystallizable fragment receptors on other cells, such as macrophages and natural killer cells (23). To overcome these challenges, antibody fragments such as the minibody, diabody, and single-domain antibody formats and other engineered protein scaffolds have been evaluated as radiotracers (24,25). These smaller vectors are likely to exhibit better tissue penetration and faster clearance and are more suited to radiolabeling with PET isotopes with shorter half-lives, making same-day imaging more feasible (26).

Radiolabeled small molecules are usually designed to target specific intracellular metabolic pathways, such as carbohydrate metabolism and DNA synthesis. The most widely used clinical PET tracer for cancer diagnosis and staging, ^{18}F -FDG, has also

TABLE 1
Candidate Biomarkers for PET Imaging of T-Cell Responses

Target	Type	Stage	Application
OX40	Surface biomarker	Preclinical	TILs, GVHD
CTLA-4	Surface biomarker	Preclinical	TILs, GVHD
CD3	Surface biomarker	Preclinical	TILs
CD4	Surface biomarker	Preclinical	TILs, IBD, HSCT, lymphoid organs
CD8	Surface biomarker	Preclinical/clinical	TILs, lymphoid organs
ICOS	Surface biomarker	Preclinical	TILs, CAR-T
PD-1	Surface biomarker	Preclinical/clinical	TILs
IL-2R	Surface biomarker	Preclinical/clinical	TILs, ONNV, lymphoid organs
TCRmu	Surface biomarker	Preclinical	CAR-T
VLA4	Surface biomarker	Preclinical	Tuberculosis
Granzyme B	Secreted biomarker	Preclinical	TILs
dCK	Intracellular enzyme	Preclinical/clinical	TILs, autoimmune hepatitis
dGK	Intracellular enzyme	Preclinical/clinical	GVHD, TILs, RA
HSV1-tk	Reporter gene/encoded protein size (46 kDa)	Preclinical/clinical	CAR-T
HSV1-sr39tk	Reporter gene/encoded protein size (42 kDa)	Preclinical	RA
PSMA	Reporter gene/encoded protein size (100 kDa)	Preclinical	CAR-T
NET	Reporter gene/encoded protein size (69 kDa)	Preclinical	TILs
SSTR2	Reporter gene/encoded protein size (41 kDa)	Preclinical	CAR-T
DHFR	Reporter gene/encoded protein size (18 kDa)	Preclinical	CAR-T
NIS	Reporter gene/encoded protein size (69 kDa)	Preclinical	CAR-T
2D12.5/G54C	Reporter gene/encoded protein size (52 kDa)	Preclinical	CAR-T

TIL = tumor infiltrating lymphocytes; CTLA-4 = cytotoxic T-lymphocyte antigen-4; IBD = inflammatory bowel disease; HSCT = hematopoietic stem cell transplantation; IL-2R = interleukin-2 receptors; ONNV = o'nyong-nyong virus; TCRmu = murine T-cell receptor β -domain; VLA4 = very late antigen-4; dCK = deoxycytidine kinase; dGK = deoxyguanosine kinase; PSMA = prostate-specific membrane antigen; NET = norepinephrine transporter; SSTR2 = somatostatin receptor 2; DHFR = dihydrofolate reductase enzyme; NIS = sodium iodide symporter; 2D12.5/G54C = antilanthanoid-DOTA antibody.

PET tracers targeting biomarkers listed here are also discussed in references 3, 22, and 35.

been explored for monitoring immune responses (27). Since the glycolytic pathway is shared by both activated immune cells and cancer cells, its specificity for the interrogation of T-cell responses is low. Rapidly proliferating T cells rely heavily on the nucleoside salvage pathway for DNA synthesis. Deoxycytidine kinase and deoxyguanosine kinase are key enzymes that regulate this pathway, motivating the development of radiolabeled small molecules targeting each. The first deoxycytidine kinase-targeting PET tracer, 1-2'-deoxy-2'-¹⁸F-fluoroarabinofuranosyl cytosine, was able to distinguish proliferating CD8-positive T cells from naïve T cells (28), but rapid catabolism in vivo impeded its clinical translation. Another deoxycytidine kinase tracer with improved metabolic stability, 2-chloro-2'-deoxy-2'-¹⁸F-fluoro-9-b-D-arabinofuranosyl-adenine, was subsequently developed and evaluated in glioblastoma patients treated with PD-1 blockade. Increased accumulation of this tracer was observed in secondary lymphoid organs and tumor tissue, and the PET signal in the latter strongly correlated with the concentration of tumor-infiltrating lymphocytes (Fig. 3C) (29). ¹⁸F-AraG, a guanosine analog with high specificity for deoxyguanosine kinase, was successfully used to detect activated T cells induced by anti-PD-1 therapy in mouse models, enabling early prediction of therapeutic response (30). Given its favorable imaging characteristics,

several clinical trials are currently evaluating the feasibility of monitoring T-cell responses with ¹⁸F-AraG (NCT04186988, NCT04726215).

Engineering cells to express reporter genes encoding proteins that can be detected with complementary PET tracers is a promising approach suited to tracking adoptively transferred T cells (31). The most extensively evaluated PET reporter gene is the herpes simplex virus type 1 thymidine kinase (*HSV1-tk*) and its mutant version *HSV1-sr39tk*. In a pilot clinical study, CD8-positive cytotoxic T lymphocytes were engineered to express both *HSV1-tk* and interleukin-13 zetakine chimeric antigen receptor (CAR) and then adoptively transferred into recurrent high-grade glioblastoma patients. ¹⁸F-FHBG was subsequently administered to monitor the trafficking and proliferation of the engineered cytotoxic T lymphocytes in the brain (Fig. 3D) (32). This landmark study demonstrated the feasibility of using the *HSV1-tk* system to clinically track engineered immune cells. The *HSV1-sr39tk* system can also act as a suicide gene on treatment with the prodrug ganciclovir, enabling ablation of CAR-T cells for safe control of potential toxicities (33). Human PET reporter genes including sodium iodide symporter, norepinephrine transporter, and somatostatin receptor 2 have also been developed to overcome the potential




FIGURE 2. PET imaging of T cells for early detection of inflammatory diseases. (A) ^{18}F -AraG enables detection of T-cell activation in cervical lymph nodes during acute GVHD (7). (B) OX40 immuno-PET allows early diagnosis of acute GVHD, before overt clinical symptoms (9). (C) ^{18}F -AraG imaging detects RA in mouse model of adjuvant-induced arthritis (11). (D) ^{89}Zr -labeled CD4-targeting cys-diabody allows detection of inflammatory bowel disease (white arrow indicating mesenteric lymph nodes) (13). CLN = cervical lymph nodes; I = intestine; M = mesenteric lymph node, S = spleen.

immunogenicity of *HSV1-tk* and evaluated for CAR-T-cell imaging. However, because of their endogenous expression, as well as the internalization of hNET and somatostatin receptor 2, their application has been limited (34). Recently, a highly promising prostate-specific membrane antigen/ ^{18}F -DCFPyL reporter system was evaluated for CD19-positive CAR-T-cell imaging (35). To specifically prevent the internalization of prostate-specific membrane antigen, human CD19-targeted CAR-T cells were transduced with an N-terminally modified variant, tPSMA^(N9del). ^{18}F -DCFPyL, a prostate-specific membrane antigen-specific PET tracer, subsequently allowed detection of these cells with high sensitivity and specificity in vivo. Importantly, quantification of the ^{18}F -DCFPyL PET signal from CD19-positive CAR-T cells in these tumors led to more accurate assessment of therapeutic response than that from measurement of CAR-T cells in peripheral blood.




FIGURE 3. PET imaging of T cells enables monitoring of treatment response in cancer immunotherapy. (A) ICOS immuno-PET with ^{89}Zr -DFO-ICOS monoclonal antibody visualizes and predicts therapeutic response in mouse model of Lewis lung cancer treated intratumorally with STING agonist and PD-1 blockade (37). (B) OX40 immuno-PET imaging enables visualization of activated T cells in A20 tumor-bearing mouse treated intratumorally with CpG (arrow represents CpG-treated tumor) (44). (C) Elevated uptake of 2-chloro-2'-deoxy-2'- ^{18}F -fluoro-9- β -D-arabino-furanosyl-adenine was observed in several lymph nodes (arrows) of recurrent glioblastoma patient after treatment with tumor lysate-pulsed DCVax and PD-1 blockade (bottom), compared with before treatment (top) (29). (D) ^{18}F -FHBG enables visualization of HSV-TK1 reporter gene-modified interleukin-13 CAR-T cells (arrows) in recurrent glioblastoma patients (top, before CAR-T infusion; bottom, after CAR-T infusion) (32). T = treated tumor.

CONCLUDING REMARKS

Here, we have reviewed state-of-the-art PET imaging approaches for in vivo T-cell visualization and their applications in T-cell-mediated diseases and cancer immunotherapy. Noninvasive PET imaging has the potential to provide comprehensive information about the distribution and abundance of immune cell subsets in real time (36,37). With these imaging approaches, physicians and research scientists can be better informed about the adaptive immune response and T-cell behaviors in different settings, which would greatly facilitate personalized medicine.

For novel T-cell PET tracers to translate from bench to bedside, lessons learned in the clinic should be brought back to the bench

to facilitate probe optimization. Although many T-cell-targeted PET tracers have yielded promising preclinical results, only a few candidates are currently under clinical investigation, and they have yet to be approved for routine clinical use. This imbalance may be attributed to redundant preclinical studies. To streamline PET tracer development, unsupervised analysis of prospective biomarkers using both preclinical and clinical “omic” datasets would help identify the optimal target for a given application. In a recent study, we used RNAseq to compare several candidate T-cell activation markers expressed on activated CD19-positive human CAR-T cells, and inducible costimulator was identified as a promising imaging biomarker that was specifically and consistently upregulated; subsequent PET imaging was focused accordingly (38). In the future, advanced techniques such as Slide-seq and CODEX (CO-Detection by indEXing) should be incorporated to identify the most relevant biomarker for T-cell imaging in a given scenario (39,40). Another reason for the imbalance is the slow adoption of novel molecular imaging techniques in the clinic. In conventional clinical workflow, anatomic imaging remains the gold standard for therapeutic response evaluation and monitoring; novel immunotherapy adjuvants are also approved by the Food and Drug Administration under certain criteria. Future clinical studies need to demonstrate that these PET approaches provide actionable insights for improved patient management. In a recent landmark study, PD-L1 imaging in cancer patients indicated that noninvasive immuno-PET outperformed RNA and immunohistochemistry-based biomarker measurements from tissue biopsies in the selection of patients most likely to benefit from PD-L1 blockade (41). These are compelling reasons to accelerate the clinical evaluation of new probes and the selection of the most promising candidates for further consideration.

To optimize the T-cell PET imaging toolbox, we should carefully compare different biomarkers, pathways, and probe formats. For example, intact antibodies exhibit higher binding affinity than small molecules but lower tissue penetration. Although the reporter gene strategy has the advantage of low background, immunogenicity usually limits its wider applications. Thus, among all the potential candidates, we should consider their affinity, specificity, sensitivity, immunogenicity, tissue penetration, and clearance and then select the best candidate for further investigation. Moreover, with current imaging probes, although we can visualize the in vivo distribution of distinct T-cell populations, we are unable to delineate their antigen specificity. A recent published study reported a novel engineered PET imaging probe—⁶⁴Cu-labeled synTac (synapse for T-cell activation)—which was able to distinguish antigen-specific CD8-positive T cells from bystander CD8-positive T cells (42). This powerful approach could allow the detection of cytotoxic CD8-positive T cells specific for invading pathogens or tumor cells, which would provide more precise assessment of cancer immunotherapy efficacy or disease severity. Finally, the groundbreaking total-body PET will substantially improve sensitivity, enabling enhanced detection of T cells while also lowering the radioactive dose typically required to acquire high-resolution images, allowing for safer repeat imaging (43).

In summary, the T-cell PET imaging toolbox has great potential for improving clinical management of cancer immunotherapy and diagnosis of T-cell-driven immunopathology. Further optimization of these approaches is still needed to overcome limitations in their specificity, sensitivity, and safety. We hope that our discussion will highlight the utility of the T-cell PET imaging toolbox to both researchers and physicians and encourage the translation and wider adoption of these tools in clinical practice.

DISCLOSURE

This work was supported by Heilongjiang Postdoctoral funding (LBH-Z18197). Israt Alam was supported by NIH/NCI (R01 CA201719-05). No other potential conflict of interest relevant to this article was reported.

ACKNOWLEDGMENTS

We thank our colleagues in the fields of molecular imaging, immunology, and oncology, among others, for their contributions toward the development of T-cell PET imaging strategies and probes discussed in this review.

REFERENCES

1. Waldman AD, Fritz JM, Lenardo MJ. A guide to cancer immunotherapy: from T cell basic science to clinical practice. *Nat Rev Immunol.* 2020;20:651–668.
2. Alcover A, Alarcón B, Di Bartolo V. Cell biology of T cell receptor expression and regulation. *Annu Rev Immunol.* 2018;36:103–125.
3. Wei W, Jiang D, Ehlerding EB, et al. Noninvasive PET imaging of T cells. *Trends Cancer.* 2018;4:359–373.
4. Pittet MJ, Grimm J, Berger CR, et al. In vivo imaging of T cell delivery to tumors after adoptive transfer therapy. *Proc Natl Acad Sci USA.* 2007;104:12457–12461.
5. Rheumatoid arthritis. *Nat Rev Dis Primers.* 2018;4:18002.
6. Narkhede M, Rybicki L, Abounader D, et al. The association of histologic grade with acute graft-versus-host disease response and outcomes. *Am J Hematol.* 2017;92:683–688.
7. Ronald JA, Kim BS, Gowrishankar G, et al. A PET imaging strategy to visualize activated T cells in acute graft-versus-host disease elicited by allogeneic hematopoietic cell transplant. *Cancer Res.* 2017;77:2893–2902.
8. Pektor S, Schlöder J, Klasen B, et al. Using immuno-PET imaging to monitor kinetics of T cell-mediated inflammation and treatment efficiency in a humanized mouse model for GvHD. *Eur J Nucl Med Mol Imaging.* 2020;47:1314–1325.
9. Alam IS, Simonetta F, Scheller L, et al. Visualization of activated T cells by OX40-ImmunoPET as a strategy for diagnosis of acute graft-versus-host disease. *Cancer Res.* 2020;80:4780–4790.
10. Narayan N, Owen DR, Taylor PC. Advances in positron emission tomography for the imaging of rheumatoid arthritis. *Rheumatology (Oxford).* 2017;56:1837–1846.
11. Franc BL, Goth S, MacKenzie J, et al. In vivo PET imaging of the activated immune environment in a small animal model of inflammatory arthritis. *Mol Imaging.* 2017;16:1536012117712638.
12. van der Krogt JMA, van Binsbergen WH, van der Laken CJ, et al. Novel positron emission tomography tracers for imaging of rheumatoid arthritis. *Autoimmun Rev.* 2021;20:102764.
13. Freise AC, Zettlitz KA, Salazar FB, et al. Immuno-PET in inflammatory bowel disease: imaging CD4-positive T cells in a murine model of colitis. *J Nucl Med.* 2018;59:980–985.
14. Hiam-Galvez KJ, Allen BM, Spitzer MH. Systemic immunity in cancer. *Nat Rev Cancer.* 2021;21:345–359.
15. Heppt MV, Steeb T, Schlager JG, et al. Immune checkpoint blockade for unresectable or metastatic uveal melanoma: a systematic review. *Cancer Treat Rev.* 2017;60:44–52.
16. Tabchi S, Kassouf E, Rassy EE, et al. Management of stage III non-small cell lung cancer. *Semin Oncol.* 2017;44:163–177.
17. Wołaczewicz M, Hryniewicz R, Grywalska E, et al. Immunotherapy in bladder cancer: current methods and future perspectives. *Cancers (Basel).* 2020;12:1181.
18. Stenger D, Stief TA, Kaeuferle T, et al. Endogenous TCR promotes in vivo persistence of CD19-CAR-T cells compared to a CRISPR/Cas9-mediated TCR knockout CAR. *Blood.* 2020;136:1407–1418.
19. Valero C, Lee M, Hoen D, et al. Response rates to anti-PD-1 immunotherapy in microsatellite-stable solid tumors with 10 or more mutations per megabase. *JAMA Oncol.* 2021;7:739–743.
20. Tazdait M, Mezquita L, Lahmar J, et al. Patterns of responses in metastatic NSCLC during PD-1 or PDL-1 inhibitor therapy: comparison of RECIST 1.1, irRECIST and iRECIST criteria. *Eur J Cancer.* 2018;88:38–47.
21. Borcoman E, Nandikolla A, Long G, et al. Patterns of response and progression to immunotherapy. *Am Soc Clin Oncol Educ Book.* 2018;38:169–178.
22. Krekorian M, Fruhwirth GO, Srinivas M, et al. Imaging of T-cells and their responses during anti-cancer immunotherapy. *Theranostics.* 2019;9:7924–7947.
23. Vivier D, Sharma SK, Adumeau P, et al. The impact of FcγRI binding on immuno-PET. *J Nucl Med.* 2019;60:1174–1182.

24. Zhao H, Wang C, Yang Y, et al. ImmunoPET imaging of human CD8⁺ T cells with novel ⁶⁸Ga-labeled nanobody companion diagnostic agents. *J Nanobiotechnology*. 2021;19:42.
25. Freise AC, Wu AM. In vivo imaging with antibodies and engineered fragments. *Mol Immunol*. 2015;67:142–152.
26. Wu AM. Engineered antibodies for molecular imaging of cancer. *Methods*. 2014;65:139–147.
27. Valentinuzzi D, Vrankar M, Boc N, et al. [¹⁸F]FDG PET immunotherapy radiomics signature (iRADIOMICS) predicts response of non-small-cell lung cancer patients treated with pembrolizumab. *Radiol Oncol*. 2020;54:285–294.
28. Radu CG, Shu CJ, Nair-Gill E, et al. Molecular imaging of lymphoid organs and immune activation by positron emission tomography with a new [¹⁸F]-labeled 2'-deoxycytidine analog. *Nat Med*. 2008;14:783–788.
29. Antonios JP, Soto H, Everson RG, et al. Detection of immune responses after immunotherapy in glioblastoma using PET and MRI. *Proc Natl Acad Sci USA*. 2017;114:10220–10225.
30. Levi J, Lam T, Goth SR, et al. Imaging of activated T cells as an early predictor of immune response to anti-PD-1 therapy. *Cancer Res*. 2019;79:3455–3465.
31. Iafrate M, Fruhwirth GO. How non-invasive in vivo cell tracking supports the development and translation of cancer immunotherapies. *Front Physiol*. 2020;11:154.
32. Keu KV, Witney TH, Yaghoubi S, et al. Reporter gene imaging of targeted T cell immunotherapy in recurrent glioma. *Sci Transl Med*. 2017;9:eaag2196.
33. Murty S, Labanieh L, Murty T, et al. PET reporter gene imaging and ganciclovir-mediated ablation of chimeric antigen receptor T cells in solid tumors. *Cancer Res*. 2020;80:4731–4740.
34. Abousaway O, Rakhshandehroo T, Van den Abbeele AD, et al. Noninvasive imaging of cancer immunotherapy. *Nanotheranostics*. 2021;5:90–112.
35. Minn I, Huss DJ, Ahn HH, et al. Imaging CAR T cell therapy with PSMA-targeted positron emission tomography. *Sci Adv*. 2019;5:eaaw5096.
36. Mayer AT, Gambhir SS. The immunoimaging toolbox. *J Nucl Med*. 2018;59:1174–1182.
37. Xiao Z, Mayer AT, Nobashi TW, et al. ICOS is an indicator of T-cell-mediated response to cancer immunotherapy. *Cancer Res*. 2020;80:3023–3032.
38. Simonetta F, Alam IS, Lohmeyer JK, et al. Molecular imaging of chimeric antigen receptor T cells by ICOS-immunoPET. *Clin Cancer Res*. 2021;27:1058–1068.
39. Rodrigues SG, Stickels RR, Goeva A, et al. Slide-seq: a scalable technology for measuring genome-wide expression at high spatial resolution. *Science*. 2019;363:1463–1467.
40. Goltsev Y, Samusik N, Kennedy-Darling J, et al. Deep profiling of mouse splenic architecture with CODEX multiplexed imaging. *Cell*. 2018;174:968–981.e15.
41. Bensch F, van der Veen EL, Lub-de Hooge MN, et al. ⁸⁹Zr-atezolizumab imaging as a non-invasive approach to assess clinical response to PD-L1 blockade in cancer. *Nat Med*. 2018;24:1852–1858.
42. Woodham AW, Zeigler SH, Zeyang EL, et al. In vivo detection of antigen-specific CD8⁺ T cells by immuno-positron emission tomography. *Nat Methods*. 2020;17:1025–1032.
43. Cherry SR, Jones T, Karp JS, et al. Total-body PET: maximizing sensitivity to create new opportunities for clinical research and patient care. *J Nucl Med*. 2018;59:3–12.
44. Alam IS, Mayer AT, Sagiv-Barfi I, et al. Imaging activated T cells predicts response to cancer vaccines. *J Clin Invest*. 2018;128:2569–2580.