

In Vivo Evaluation of 6 Analogs of ^{11}C -ER176 as Candidate ^{18}F -Labeled Radioligands for 18-kDa Translocator Protein

Jae-Hoon Lee^{1,2}, Fabrice G. Siméon¹, Jehi-San Liow¹, Cheryl L. Morse¹, Robert L. Gladding¹, Jose A. Montero Santamaria¹, Ioline D. Henter¹, Sami S. Zoghbi¹, Victor W. Pike¹, and Robert B. Innis¹

¹Molecular Imaging Branch, National Institute of Mental Health, National Institutes of Health, Bethesda, Maryland; and ²Department of Nuclear Medicine, Yonsei University College of Medicine, Seoul, South Korea

Because of its excellent ratio of specific to nondisplaceable uptake, the radioligand ^{11}C -ER176 can successfully image 18-kDa translocator protein (TSPO), a biomarker of inflammation, in the human brain and accurately quantify target density in homozygous low-affinity binders. Our laboratory sought to develop an ^{18}F -labeled TSPO PET radioligand based on ER176 with the potential for broader distribution. This study used generic ^{11}C labeling and in vivo performance in the monkey brain to select the most promising among 6 fluorine-containing analogs of ER176 for subsequent labeling with longer-lived ^{18}F . **Methods:** Six fluorine-containing analogs of ER176—3 fluoro and 3 trifluoromethyl isomers—were synthesized and labeled by ^{11}C methylation at the secondary amide group of the respective *N*-desmethyl precursor. PET imaging of the monkey brain was performed at baseline and after blockade by *N*-butan-2-yl-1-(2-chlorophenyl)-*N*-methylisoquinoline-3-carboxamide (PK11195). Uptake was quantified using radiometabolite-corrected arterial input function. The 6 candidate radioligands were ranked for performance on the basis of 2 in vivo criteria: the ratio of specific to nondisplaceable uptake (i.e., nondisplaceable binding potential [BP_{ND}]) and the time stability of total distribution volume (V_{T}), an indirect measure of lack of radiometabolite accumulation in the brain. **Results:** Total TSPO binding was quantified as V_{T} corrected for plasma free fraction ($V_{\text{T}}/f_{\text{p}}$) using Logan graphical analysis for all 6 radioligands. $V_{\text{T}}/f_{\text{p}}$ was generally high at baseline ($222 \pm 178 \text{ mL}\cdot\text{cm}^{-3}$) and decreased by 70%–90% after preblocking with PK11195. BP_{ND} calculated using the Lassen plot was 9.6 ± 3.8 ; the *o*-fluoro radioligand exhibited the highest BP_{ND} (12.1), followed by the *m*-trifluoromethyl (11.7) and *m*-fluoro (8.1) radioligands. For all 6 radioligands, V_{T} reached 90% of the terminal 120-min values by 70 min and remained relatively stable thereafter, with excellent identifiability ($\text{SEs} < 5\%$), suggesting that no significant radiometabolites accumulated in the brain. **Conclusion:** All 6 radioligands had good BP_{ND} and good time stability of V_{T} . Among them, the *o*-fluoro, *m*-trifluoromethyl, and *m*-fluoro compounds were the 3 best candidates for development as radioligands with an ^{18}F label.

Key Words: translocator protein; neuroinflammation; PET; specific-to-nondisplaceable uptake; radiometabolites

J Nucl Med 2021; 63:1252–1258
DOI: 10.2967/jnumed.121.263168

Received Sep. 8, 2021; revision accepted Nov. 30, 2021.
For correspondence or reprints, contact Jae-Hoon Lee (jae-hoon.lee@nih.gov).
Published online Jan. 13, 2022.
COPYRIGHT © 2022 by the Society of Nuclear Medicine and Molecular Imaging.

The mitochondrial protein 18-kDa translocator protein (TSPO) is highly expressed in phagocytic inflammatory cells, including activated microglia and reactive astrocytes in the brain and macrophages in the periphery (1,2). Although numerous PET radioligands have been developed to image TSPO, several limitations have restricted their clinical utility for quantifying inflammation in the brain. Of these radioligands, the first-generation TSPO radioligand ^{11}C -(*R*)-*N*-butan-2-yl-1-(2-chlorophenyl)-*N*-methylisoquinoline-3-carboxamide (PK11195) has been the most extensively studied. ^{11}C -(*R*)-PK11195 has high affinity for TSPO; however, its utility is limited by a low ratio of specific to nondisplaceable uptake (i.e., nondisplaceable binding potential [BP_{ND}]) in the brain and by the relatively short half-life of ^{11}C (20 min) (3,4). Second-generation radioligands, such as ^{11}C -PBR28, offer a higher in vivo TSPO-specific signal but suffer from sensitivity to the single-nucleotide polymorphism *rs6971* (5,6), such that low-affinity binders (LABs) have too little TSPO binding to be accurately measured. This sensitivity to the single-nucleotide polymorphism both complicates the interpretation of results and requires genotyping to exclude LABs before imaging. Consequently, new and more effective radioligands are needed to image TSPO.

Third-generation TSPO radioligands were designed to have adequately high BP_{ND} across all *rs6971* genotypes for reliable quantification and to lack the brain radiometabolite accumulation that interferes with the low specific signal in LABs. In this context, ^{11}C -ER176 is arguably the most promising third-generation TSPO radioligand for clinical research (2,7). More specifically, it displays high specific binding (>80%), an adequately high BP_{ND} in LABs, and good time stability of total distribution volume (V_{T}) across all genotypes (8,9). The whole-brain BP_{ND} of ^{11}C -ER176 for LABs was 1.4 ± 0.8 , which is about the same as that for high-affinity binders with ^{11}C -PBR28 (~1.2). For all 3 genotypes, the V_{T} values of ^{11}C -ER176 stabilized by 60–90 min to within 10% of their final 120-min values, suggesting that no significant amount of radiometabolites accumulated in the brain (8).

In contrast to ^{11}C , ^{18}F has several benefits for clinical PET imaging. Its relatively longer half-life (110 vs. 20 min) allows imaging for a longer period and for broader distribution of the radioligand from a central radiopharmacy, giving ^{18}F -labeled radioligands greater flexibility for widespread use. However, the structure of ER176 does not contain a fluorine atom for labeling with ^{18}F . Thus, as a first step toward developing an ^{18}F -labeled third-generation radioligand, our laboratory synthesized 6 new fluorine-containing analogs of ER176—3 isomers with a fluoro group and 3 with a trifluoromethyl group at each of 3 positions (*ortho*, *meta*, and *para*) of the pendant aryl ring (Fig. 1) (10). In vitro studies demonstrated that all 6 analogs had high affinity for human TSPO (binding affinity, 1.2–7.0 nM) and could




FIGURE 1. Chemical structures of ^{11}C -ER176 and 6 fluorine-containing analogs. K_i = binding affinity.

be successfully labeled with ^{11}C with good yield (66%–81%, decay-corrected) and excellent chemical (>95%) and radiochemical (>99%) purities. Selecting the most promising radioligand among the 6 candidates to label with ^{18}F would significantly reduce the time and cost of future investigations; however, *in vitro* data alone would not provide sufficient guidance for this decision.

The present study used *in vivo* performance in the monkey brain to select the most promising of the 6 ^{11}C -labeled analogs of ER176 for subsequent ^{18}F -radiolabeling. The 2 primary performance criteria were BP_{ND} and the stability of V_T over time, which is an indirect measure of lack of radiometabolite accumulation in the brain. Both baseline and preblocked PET scans in the monkey brain were obtained, and brain uptake was quantified as V_T using the radiometabolite-corrected arterial input function.

MATERIALS AND METHODS

Radiochemistry

^{11}C -ER176 was synthesized as previously described (11), with a molar activity of $106 \pm 65 \text{ GBq}/\mu\text{mol}$ ($n = 4$) at the time of injection and radiochemical purity of $97.1\% \pm 4.4\%$.

Six fluorine-containing analogs of ER176 (Fig. 1) were labeled with ^{11}C at the tertiary amide group by ^{11}C -methylation of the respective *N*-desmethyl precursor as previously described (10). Briefly, the labeling precursors were synthesized by amidation of 4-oxo-3*H*-quinazoline-2-carboxylic acid followed by palladium-catalyzed coupling with appropriate fluorophenylboronic acids for the fluoro isomers—*o*-fluoro (SF12063), *m*-fluoro (SF12051), and *p*-fluoro (SF12052)—and trifluoromethylphenylboronic acids for the trifluoromethyl isomers—*o*-trifluoromethyl (SF12050), *m*-trifluoromethyl (SF12057), and *p*-trifluoromethyl (SF12054). Six ^{11}C -labeled TSPO radioligands were then obtained by methylation at the secondary amide group of the respective *N*-desmethyl precursors in dimethyl sulfoxide with ^{11}C -methyl iodide. The molar activity was $222 \pm 162 \text{ GBq}/\mu\text{mol}$ ($n = 18$) for the fluoro radioligands and $111 \pm 71 \text{ GBq}/\mu\text{mol}$ ($n = 10$) for the trifluoromethyl radioligands at the time of injection (Supplemental Table 1; supplemental materials are available at <http://jnm.snmjournals.org>). The radiochemical purity of all radioligands was $99.6\% \pm 0.5\%$ ($n = 28$).

Animals

In vivo experiments were performed on 9 healthy male rhesus monkeys (body weight, $12.4 \pm 1.3 \text{ kg}$). Anesthesia was maintained with 1%–2% isoflurane and 98% O_2 for the duration of the study. The head was firmly fixed by gauze and tape to the camera bed holder. Body temperature was maintained with air blankets; temperature, oxygen saturation, blood pressure, and end-tidal CO_2 were monitored for the duration of the study. All animal studies were conducted in compliance with the *Guide for the Care and Use of Laboratory Animals* (12) and were approved by the National Institute of Mental Health Animal Care and Use Committee.

PET Data Acquisition

The baseline PET scan was acquired for all radioligands (injected activity, $249 \pm 78 \text{ MBq}$) using a microPET Focus 220 scanner (Siemens Medical Solutions), with a frame duration ranging from 30 s to 10 min. For the preblocked scans, racemic PK11195 (5 mg/kg) was intravenously administered 5–10 min before the radioligand. All PET scans were acquired for 120 min at baseline and 90 min after preblocking. Concurrent arterial blood sampling was performed in all scans to obtain a radiometabolite-corrected input function for quantification. PET images were reconstructed using a Fourier rebinning algorithm plus 2-dimensional filtered backprojection with attenuation and scatter correction.

Measurement of Parent Radioactivity in Plasma

Fifteen blood samples were drawn from an implanted port in the femoral artery during the PET scan every 15 s for the first 2 min, followed by sampling at 3, 5, 10, 30, 60, 90, and 120 min (varying from 1.0 to 3.0 mL); 14 samples were drawn during the 90-min PET scan. The parent radioligand was separated from radiometabolites as previously described (13). Plasma parent and whole-blood activity concentration were fitted with a triexponential function. The plasma free fraction (f_p) was measured by ultrafiltration as previously described (14).

Kinetic Analysis

All kinetic analyses were performed using PMOD, version 3.9 (PMOD Technologies Ltd.). V_T was estimated with Logan graphical analysis (LGA) using 90 min of brain time-activity curves and the radiometabolite-corrected arterial input function for both baseline and preblocked studies. Details on the kinetic analysis are provided in the supplemental materials (15).

Estimating Ratio of Specific to Nondisplaceable Uptake

BP_{ND} was used to compare the performance of the 6 radioligands; this measure is a ratio of receptor-specific uptake (V_S ; $V_T - V_{\text{ND}}$) to nondisplaceable uptake (V_{ND} ; free plus nonspecific binding) at equilibrium. In our opinion, BP_{ND} is a better binding measure than V_T because BP_{ND} directly quantifies specific binding. BP_{ND} is also more suitable than V_S because it is a signal-to-noise measurement that takes background into consideration. BP_{ND} was calculated as follows (16):

$$BP_{\text{ND}} = \frac{V_T - V_{\text{ND}}}{V_{\text{ND}}} = \frac{V_T}{V_{\text{ND}}} - 1,$$

where V_T refers to V_T at baseline and V_{ND} was estimated by the Lassen plot.

Because only unbound parent radioligand contributes to specific binding to the receptor, and because f_p was significantly different before and

after preblocking, brain uptake and BP_{ND} were calculated and compared for the 6 radioligands using V_T and V_{ND} corrected for f_p as V_T/f_p and V_{ND}/f_p , respectively.

Time Stability Analysis of V_T

To determine the minimal scan duration needed to reliably measure V_T and to indirectly assess whether radiometabolites enter the brain, time stability was evaluated using 120 min of baseline PET data with a truncated acquisition duration from 30 to 120 min in 10-min increments. The identifiability of V_T was also evaluated as percentage SE at each truncated scan duration.

Ex Vivo Blood Cell Analysis

The utility of radioligand uptake was investigated in blood cells as a surrogate for brain uptake. The radioactivity concentration in ex vivo blood cells was calculated both at baseline and in preblocked studies as follows (17):

$$C_{BC} = C_{PL} + \frac{(C_{WB} - C_{PL})}{\text{hematocrit}}$$

Here, C_{BC} , C_{PL} , and C_{WB} indicate radioactivity concentrations in the blood cells, plasma, and whole blood, respectively. The distribution volume in blood cells (V_{BC}) was then obtained by dividing the radioactivity concentrations in the blood cells by the radioactivity concentration of the parent radioligand in plasma and then corrected for f_p (V_{BC}/f_p). The correlation between V_{BC}/f_p and whole-brain V_T/f_p and between percentage blockade calculated using V_{BC}/f_p and whole-brain V_T/f_p was assessed using linear regression analysis.

Replication Study and Statistical Analysis

Based on the results of the first scans, replication studies were conducted for selected radioligands in different monkeys to confirm either the radioligands' most favorable properties or aberrant data. When multiple experiments were performed with the same radioligand, quantitative results are presented as mean \pm SD. Statistical significance

was set at a P value of less than 0.05. All statistical analyses were conducted with Prism, version 5 (GraphPad Software).

RESULTS

Uptake in Monkey Brain

Brain radioactivity at baseline increased rapidly and peaked (SUV_{peak} , 2.7) at about 25 min after injection (Figs. 2A and 2B) except for the *p*-trifluoromethyl radioligand, which showed very slow uptake that peaked at 75 min. The *m*-trifluoromethyl radioligand showed the highest SUV_{peak} (3.4), and the *m*-fluoro radioligand showed the lowest SUV_{peak} (2.6). The brain region with the highest SUV_{peak} was the striatum (3.9), followed by the thalamus (3.2) and cerebellum (3.1); the parietal cortex had the lowest SUV_{peak} (2.4). Peak radioactivity uptake in all regions was followed by a smooth decrease in radioactivity level; for example, radioactivity in the whole brain decreased by 37% at 90 min after injection. In preblocked scans, all 6 radioligands showed a similar brain uptake pattern. Brain radioactivity rapidly increased and reached an SUV_{peak} of 4.1 at 3.5 min after injection, followed by a rapid decline and then a slow washout (Figs. 2C and 2D).

Plasma Concentration of Parent Radioligand

The concentration of parent radioligands in plasma peaked at 1.0–1.3 min after injection at baseline and then rapidly declined, followed by a slow terminal clearance phase. The fitting of plasma parent curves converged by triexponential function in all experiments (Fig. 3). The plasma parent fraction at baseline, expressed as a percentage of total plasma radioactivity, declined rapidly and reached 50% at 25.6 ± 11.6 min for the *m*-fluoro, *p*-fluoro, and *o*-trifluoromethyl radioligands (Supplemental Fig. 1). In contrast, the *o*-fluoro and *m*-trifluoromethyl radioligands showed a slower decline, reaching 50% at 50.7 ± 14.5 min after injection, and the *p*-trifluoromethyl radioligand exhibited an unusually high plasma parent fraction (>90%) during the entire 120 min of the scan. In preblocked scans,

the parent radioactivity concentrations in plasma showed a rapid increase, peaking at 1.0–1.5 min, followed by a fast washout and then a slow terminal clearance phase in all radioligands. As in our prior studies (11,18), the peak concentration of parent radioligand in plasma was much higher in the preblocked scans than in the baseline scans, because PK11195 blocks the distribution of radioligand to peripheral organs—such as the lung and kidneys—that have high densities of TSPO. The temporal changes in parent fraction in plasma were similar for all radioligands, characterized by a rapid decline that reached less than 50% by 30 min, with a subsequent gradual decline.

Reversed-phase high-performance liquid chromatography of plasma revealed at least 5 radiometabolites, all of which appeared less lipophilic than the parent radioligand. A lipophilic radiometabolite appeared in plasma from all radioligand experiments, except the baseline study of the *p*-trifluoromethyl radioligand. Nonetheless, the amount was negligible ($0.07\% \pm 0.08\%$ of total plasma radioactivity across all arterial samples). The f_p was $17.3\% \pm 8.2\%$ at baseline for the




FIGURE 2. Time-activity curves of whole-brain uptake in baseline and preblocked scans for fluoro (A and C) and trifluoromethyl (B and D) radioligands. Points represent $SUV_{mean} (\pm SD)$. Conc = concentration.




FIGURE 3. Time course of radioactivity concentrations in plasma at baseline and in preblocked scans for fluoro (A and C) and trifluoromethyl (B and D) radioligands. Points represent $SUV_{mean} (\pm SD)$. Conc = concentration.

fluoro radioligands, whereas it was significantly higher ($22.3\% \pm 11.3\%$) after preblocking ($n = 9, P = 0.017$). In contrast, there was no significant difference in f_p at baseline and after preblocking for the trifluoromethyl radioligands; f_p was $10.1\% \pm 2.9\%$ at baseline and $10.6\% \pm 3.1\%$ at preblock ($n = 5, P = 0.423$).

highest ratio of specific to nondisplaceable uptake (BP_{ND} , 12.1), followed by the *m*-trifluoromethyl (11.7) and *m*-fluoro (8.1) radioligands (Table 1). The Lassen plot analysis displayed excellent linear correlations ($R^2 > 0.90$) at high receptor occupancies ($>87\%$) in all radioligands (Supplemental Fig. 2). Estimated V_{ND}/f_p ($mL \cdot cm^{-3}$)

Kinetic Analysis

Brain uptake was well quantified by LGA, which does not require specific compartment configurations. Regional brain uptake was reliably quantified as V_T with excellent identifiability ($SE < 10\%$) in all baseline and preblocked studies. The trifluoromethyl radioligands generally showed higher V_T/f_p ($mL \cdot cm^{-3}$) for baseline and preblocked conditions than the fluoro radioligands (363 vs. 138, respectively, at baseline and 50 vs. 23, respectively, at preblock) (Fig. 4). The *p*-trifluoromethyl radioligand had the highest V_T/f_p (493) at baseline, followed by the *m*-trifluoromethyl (411) and *o*-fluoro (223) radioligands (Table 1). The 3 brain regions with the highest V_T/f_p were the striatum (286), thalamus (283), and frontal cortex (269). The 3 with the lowest V_T/f_p were the cerebellum (208), amygdala (228), and occipital cortex (243). The blockade by PK11195 was $82.5\% \pm 7.3\%$ and was similar for all 6 radioligands.

Estimating Ratio of Specific to Nondisplaceable Uptake

The *o*-fluoro radioligand exhibited the highest ratio of specific to nondisplaceable uptake (BP_{ND} , 12.1), followed by the *m*-trifluoromethyl (11.7) and *m*-fluoro (8.1) radioligands (Table 1). The Lassen plot analysis displayed excellent linear correlations ($R^2 > 0.90$) at high receptor occupancies ($>87\%$) in all radioligands (Supplemental Fig. 2). Estimated V_{ND}/f_p ($mL \cdot cm^{-3}$)




FIGURE 4. Average parametric images of total TSPO binding (V_T/f_p) for ^{11}C -ER176 and 6 analogs in monkey brain at baseline (top row) and preblocked scans (bottom row). Each V_T/f_p image was generated using 0–90 min of PET data obtained via LGA.

TABLE 1

Comparison of V_T/f_P , Occupancy, V_{ND}/f_P , and BP_{ND} of Whole Brain Among ^{11}C -ER176 and 6 Fluorine-Containing Analogs

Compound	V_T/f_P ($\text{mL}\cdot\text{cm}^{-3}$)			Occupancy (%)	V_{ND}/f_P ($\text{mL}\cdot\text{cm}^{-3}$)	BP_{ND}	Binding affinity ratio*
	Baseline	Preblock	Blockade (%)				
Reference (^{11}C -ER176)	185.8	21.3	88.5	98.6	18.6	8.9	1.3
Fluoro							
<i>o</i> -fluoro	223.8	33.3	85.0	92.9	16.6	12.1	2.8
<i>m</i> -fluoro	92.2	12.7	84.9	95.6	9.8	8.1	2.7
<i>p</i> -fluoro	45.3	13.4	70.3	89.0	7.9	4.8	2.9
Trifluoromethyl							
<i>o</i> -trifluoromethyl	119.7	31.5	73.5	87.1	18.5	5.5	5.5
<i>m</i> -trifluoromethyl	411.0	47.6	86.9	95.0	32.6	11.7	2.0
<i>p</i> -trifluoromethyl	493.0	73.6	84.8	97.7	63.6	6.7	0.8

*Ratio of binding affinity (in nM) in LABs to that in high-affinity binders. Data are mean.

ranged mainly between 5.0 and 35.0, except for the *p*-trifluoromethyl radioligand (63.6). V_{ND}/f_P was smallest for the *p*-fluoro radioligands (7.9), followed by the *m*-fluoro (9.8) and *o*-fluoro (16.6) radioligands (Table 1). As shown in Supplemental Table 2, the results were similar to the analysis without f_P correction.

Time Stability of V_T

Whole-brain V_T asymptotically reached terminal values and converged within 10% of terminal values by the 70th minute of a 120-min scan (Figs. 5A and 5B). V_T remained almost stable for the last 50 min, showing an average change of 4.7%, and could be quantified with excellent identifiability ($\text{SE} < 5\%$) (Supplemental Fig. 3). The fluoro and trifluoromethyl radioligands took a similar time to achieve a stable V_T (70 min). The relatively stable V_T measurement over the 70–120 min of the scan suggested no significant accumulation of radiometabolites in the brain.

Performance Comparison with ^{11}C -ER176

o-fluoro, *m*-trifluoromethyl, and *m*-fluoro radioligands showed a BP_{ND} similar to or higher than that of ^{11}C -ER176 (8.9), whereas the other 3 showed slightly lower values (Table 1). Regarding the time stability of V_T , it took a similar time for the whole-brain

V_T values of ^{11}C -ER176 and its 6 fluorine-containing analogs to reach and remain stable within 10% of their terminal values (90 vs. 70 min) (Fig. 5; Supplemental Fig. 4).

Correlation Between Radioligand Uptake in Blood Cells and Brain

Ex vivo blood cell uptake (V_{BC}/f_P) correlated well with in vivo whole-brain V_T/f_P in both baseline ($R^2 = 0.938$, $P < 0.001$) and preblocked ($R^2 = 0.750$, $P = 0.012$) studies (Supplemental Fig. 5). The percentage blockade of V_{BC}/f_P by PK11195 also correlated significantly with that of whole-brain V_T/f_P ($R^2 = 0.583$, $P = 0.046$).

DISCUSSION

Of the 6 ^{11}C -ER176 analogs developed by our laboratory, the present study found that the *o*-fluoro, *m*-trifluoromethyl, and *m*-fluoro compounds were the most promising. All 3 had a high BP_{ND} (the ratio of specific to nondisplaceable uptake) and a stable V_T measured over time in the monkey brain, consistent with the lack of radiometabolite accumulation. Specifically, the *o*-fluoro radioligand had the third-highest V_T/f_P at baseline (120 $\text{mL}\cdot\text{cm}^{-3}$), 96% of which was specifically bound to TSPO, and the highest BP_{ND} (12.1), making it potentially the most promising of all 6 ^{11}C -ER176 analogs. The *m*-trifluoromethyl radioligand showed the second-highest V_T/f_P at baseline (411 $\text{mL}\cdot\text{cm}^{-3}$), high specific binding to TSPO (96%), and high BP_{ND} (11.7). The *m*-fluoro radioligand also showed a high V_T/f_P at baseline (92 $\text{mL}\cdot\text{cm}^{-3}$), high specific binding to TSPO (95.6%), and a high BP_{ND} (8.1). For these 3 radioligands, V_T reached 90% of terminal 120-min values by 70 min and remained relatively stable thereafter, with excellent identifiability ($\% \text{SE} < 5\%$), suggesting that no significant radiometabolites accumulated in the brain.

In the present study, LGA was used to quantify and compare brain uptake across




FIGURE 5. Time stability analysis of whole-brain V_T for fluoro (A) and trifluoromethyl (B) radioligands. V_T was calculated via LGA and normalized to terminal V_T at 120 min. Points represent mean normalized V_T ($\pm \text{SD}$).

all 6 radioligands because neither the 1- nor the 2-tissue-compartment models fitted perfectly for all studies. With LGA, V_T was quantified with excellent identifiability ($SE < 10\%$) in all baseline and preblocked studies. Logan-derived V_{ND}/f_p and BP_{ND} were within acceptable ranges and became less variable within each of the fluoro and trifluoromethyl radioligands. However, LGA tends to underestimate V_T and BP_{ND} , typically by 10%–20% depending on both the noise level and the radioligand concentration (19,20). The degree of underestimation in this study remains uncertain; however, given the same target (TSPO) and similar performance measures, we believe that the underestimation, if any, would have been consistent across all 6 radioligands and would have led to the same finding—that is, that the *o*-fluoro, *m*-trifluoromethyl, and *m*-fluoro analogs were the 3 best candidates.

Evaluation of V_T stability with respect to scan duration allows one to determine the minimum scan time required for obtaining a stable V_T and to indirectly check the possibility that radiometabolites are accumulating in the brain. Acceptable scan durations typically occur when V_T in all regions approaches 10% of a terminal value. If brain-penetrant radiometabolites accumulate in the brain, V_T is expected to increase continuously, making the values unlikely to plateau and stabilize. Whether radiometabolites enter and accumulate in the brain is critical for imaging human TSPO, especially in LABs who have low specific binding. Although some small differences were noted between the radioligands, the present study found that, for all 6 radioligands, whole-brain V_T at baseline reached 10% of the terminal value by 70 min and 5% by 90 min. This similarity in time stability suggests that radiometabolites are unlikely to have interfered with brain uptake.

Interestingly, the present study found that ex vivo blood cells were a useful surrogate for brain tissue. V_{BC} at baseline and under preblocked conditions and the percentage blockade by PK11195 correlated well with those of whole-brain V_T/f_p . The similarity between the 2 organs can be used to evaluate radioligands that specifically bind to receptors expressed in both the brain and blood cells. For example, the blood cell analysis could be used as a quick screening tool for candidate radioligands or a nonimaging supplement to validate imaging results, thus aiding the development of new PET radioligands. However, the utility of ex vivo blood cells should be verified for individual radioligands because the 2 organs differ in their efflux systems and compartmental configurations.

One of the advantages of ^{11}C -ER176 is that it has adequately high TSPO-specific binding for quantification across all *rs6971* genotypes. A previous postmortem analysis of human brain tissue measured the in vitro binding affinities to all *rs6971* genotypes for ER176 and these 6 fluorine-containing analogs (8,10). As summarized in Table 1, the *p*-trifluoromethyl compound showed the smallest ratio (i.e., binding affinity difference) (0.8) between high-affinity binders and LABs, followed by ER176 (1.3), the *m*-trifluoromethyl (2.0) compound, the 3 fluoro compounds (2.7–2.9), and the *o*-trifluoromethyl (5.4) compound. Building on this finding, all 6 analogs investigated here are expected to achieve an adequately high specific signal in LABs, at least higher than PBR28 (55.0) (5), which has been the most widely used of the second-generation TSPO radioligands.

Taken together, these in vitro and in vivo data suggest that the *o*-fluoro, *m*-trifluoromethyl, and *m*-fluoro compounds appear to be the most promising analogs. However, all 6 radioligands performed well and with only small interligand differences (Table 1). For example, the 3 fluoro radioligands exhibited similar BP_{ND}

(4.8–12.1), time stability of V_T (all achieved stable V_T by 70 min), and binding affinity ratios (2.7–2.9), and all these measures were comparable to those of ^{11}C -ER176. Nevertheless, unexpected labeling issues (e.g., unsatisfactory yield, or perhaps only marginally acceptable molar activity for labeling trifluoromethyl groups) for ^{18}F radioligands may also arise in their development, limiting the use of particular radioligands despite solid performance in their ^{11}C -labeled forms. Thus, though the candidate radioligands have now been ranked, priorities for subsequent ^{18}F radiolabeling will also be affected by the time and effort needed for radiochemistry. Moving forward, none of the 6 compounds will be excluded.

The underlying assumption of this study is that ^{18}F -labeled radioligands will perform similarly to their ^{11}C -labeled versions. However, the performance of an ^{11}C -labeled radioligand does not always guarantee success in its ^{18}F -labeled form. ^{18}F labeling at a position different from an ^{11}C position in the molecular structure may change the spectrum of radiometabolites and generate unexpected brain-penetrating radiometabolites. Defluorination may also occur to confound the accurate quantification of receptor-specific brain uptake. Although strategic positioning of an ^{18}F label can reduce the production of these troublesome radiometabolites, the in vivo metabolism of a radioligand (e.g., the sites of metabolic cleavage) is not always predictable and further varies across species (21).

CONCLUSION

The 6 fluorine-containing analogs of ER176 were relatively easily labeled with ^{11}C . PET imaging in the monkey brain showed that all 6 ^{11}C -labeled analogs had a good BP_{ND} and good time stability of V_T . Of the 6 ligands, the *o*-fluoro, *m*-trifluoromethyl, and *m*-fluoro compounds were arguably the 3 best candidates to radiolabel with ^{18}F , a process that is expected to be quite challenging.

DISCLOSURE

This study was funded by the Intramural Research Program of the National Institute of Mental Health, National Institutes of Health (ZIAMH002795 and ZIAMH002793). No potential conflict of interest relevant to this article was reported.

ACKNOWLEDGMENTS

We thank the dedicated staff of the Molecular Imaging Branch of the NIMH, the PET Department of the NIH Clinical Center, and the NIMH veterinary staff for help in completing the studies.

KEY POINTS

QUESTION: Which are the most promising fluorine-containing analogs of ^{11}C -ER176 for subsequent ^{18}F radiolabeling?

PERTINENT FINDINGS: This study used generic ^{11}C labeling and in vivo performance in the monkey brain— BP_{ND} and time stability of V_T —to select the most promising analogs among 6 candidates. The *o*-fluoro, *m*-trifluoromethyl, and *m*-fluoro compounds were arguably the 3 best candidates because they showed the 3 highest BP_{ND} values as well as good time stability of V_T .

IMPLICATIONS FOR PATIENT CARE: The development of an ^{18}F -labeled radioligand based on ^{11}C -ER176 would allow greater flexibility and more widespread use of TSPO PET.

REFERENCES

1. Papadopoulos V, Baraldi M, Guilarte TR, et al. Translocator protein (18kDa): new nomenclature for the peripheral-type benzodiazepine receptor based on its structure and molecular function. *Trends Pharmacol Sci.* 2006;27:402–409.
2. Meyer JH, Cervenka S, Kim MJ, Kreisl WC, Henter ID, Innis RB. Neuroinflammation in psychiatric disorders: PET imaging and promising new targets. *Lancet Psychiatry.* 2020;7:1064–1074.
3. Chauveau F, Boutin H, Van Camp N, Dolle F, Tavitian B. Nuclear imaging of neuroinflammation: a comprehensive review of [¹¹C]PK11195 challengers. *Eur J Nucl Med Mol Imaging.* 2008;35:2304–2319.
4. Le Fur G, Perrier ML, Vaucher N, et al. Peripheral benzodiazepine binding sites: effect of PK 11195, 1-(2-chlorophenyl)-N-methyl-N-(1-methylpropyl)-3-isoquinolinecarboxamide. I. In vitro studies. *Life Sci.* 1983;32:1839–1847.
5. Kreisl WC, Jenko KJ, Hines CS, et al. A genetic polymorphism for translocator protein 18 kDa affects both in vitro and in vivo radioligand binding in human brain to this putative biomarker of neuroinflammation. *J Cereb Blood Flow Metab.* 2013;33:53–58.
6. Owen DR, Yeo AJ, Gunn RN, et al. An 18-kDa translocator protein (TSPO) polymorphism explains differences in binding affinity of the PET radioligand PBR28. *J Cereb Blood Flow Metab.* 2012;32:1–5.
7. Kreisl WC, Kim MJ, Coughlin JM, Henter ID, Owen DR, Innis RB. PET imaging of neuroinflammation in neurological disorders. *Lancet Neurol.* 2020;19:940–950.
8. Ikawa M, Lohith TG, Shrestha S, et al. ¹¹C-ER176, a radioligand for 18-kDa translocator protein, has adequate sensitivity to robustly image all three affinity genotypes in human brain. *J Nucl Med.* 2017;58:320–325.
9. Fujita M, Kobayashi M, Ikawa M, et al. Comparison of four ¹¹C-labeled PET ligands to quantify translocator protein 18 kDa (TSPO) in human brain: (*R*)-PK11195, PBR28, DPA-713, and ER176-based on recent publications that measured specific-to-non-displaceable ratios. *EJNMMI Res.* 2017;7:84.
10. Siméon FG, Lee JH, Morse CL, et al. Synthesis and screening in mice of fluorine-containing PET radioligands for TSPO: discovery of a promising ¹⁸F-labeled ligand. *J Med Chem.* 2021;64:16731–16745.
11. Zanotti-Fregonara P, Zhang Y, Jenko KJ, et al. Synthesis and evaluation of translocator 18 kDa protein (TSPO) positron emission tomography (PET) radioligands with low binding sensitivity to human single nucleotide polymorphism rs6971. *ACS Chem Neurosci.* 2014;5:963–971.
12. *Guide for the Care and Use of Laboratory Animals.* 8th ed. National Academy Press; 2011.
13. Zoghbi SS, Shetty HU, Ichise M, et al. PET imaging of the dopamine transporter with ¹⁸F-FECNT: a polar radiometabolite confounds brain radioligand measurements. *J Nucl Med.* 2006;47:520–527.
14. Gandelman MS, Baldwin RM, Zoghbi SS, Zea-Ponce Y, Innis RB. Evaluation of ultrafiltration for the free-fraction determination of single photon emission computed tomography (SPECT) radiotracers: beta-CIT, IBF, and iomazenil. *J Pharm Sci.* 1994;83:1014–1019.
15. Yasuno F, Brown AK, Zoghbi SS, et al. The PET radioligand [¹¹C]MePPEP binds reversibly and with high specific signal to cannabinoid CB1 receptors in nonhuman primate brain. *Neuropsychopharmacology.* 2008;33:259–269.
16. Innis RB, Cunningham VJ, Delforge J, et al. Consensus nomenclature for in vivo imaging of reversibly binding radioligands. *J Cereb Blood Flow Metab.* 2007;27:1533–1539.
17. Kanegawa N, Collste K, Forsberg A, et al. In vivo evidence of a functional association between immune cells in blood and brain in healthy human subjects. *Brain Behav Immun.* 2016;54:149–157.
18. Briard E, Zoghbi SS, Imaizumi M, et al. Synthesis and evaluation in monkey of two sensitive ¹¹C-labeled aryloxyanilide ligands for imaging brain peripheral benzodiazepine receptors in vivo. *J Med Chem.* 2008;51:17–30.
19. Logan J, Fowler JS, Volkow ND, Ding YS, Wang GJ, Alexoff DL. A strategy for removing the bias in the graphical analysis method. *J Cereb Blood Flow Metab.* 2001;21:307–320.
20. Slifstein M, Laruelle M. Effects of statistical noise on graphic analysis of PET neuroreceptor studies. *J Nucl Med.* 2000;41:2083–2088.
21. Pike VW. PET radiotracers: crossing the blood-brain barrier and surviving metabolism. *Trends Pharmacol Sci.* 2009;30:431–440.