

SUPPLEMENTAL TABLE 1. Distribution and Extent of Reduced Tracer Uptake in Patients with ARVC in Relation to the Control Group. ARVC arrhythmogenic right ventricular cardiomyopathy; MIBG ^{123}I -meta-iodo-benzylguanidine single photon emission computed tomography (SPECT); and n number. * Polar map area is displayed in relation to all 33 polar map segments (=100%). The single apical segment was not included in the analysis.

	All	ARVC		P-Value
	(n=42)	MIBG_{abnormal}	MIBG_{normal}	
		(n=25)	(n=17)	
Total number of segments, n	1386	825	561	
Total number of segments with reduced tracer uptake				
Total number of segments, n (%)	358 (25.8)	299 (83.5)	59 (16.5)	<0.0005
Septal segments, n (%)	113 (31.6)	94 (31.4)	19 (32.2)	<0.0005
Inferior segments, n (%)	111 (31.0)	95 (31.8)	16 (27.1)	<0.0005
Lateral segments, n (%)	26 (7.3)	23 (7.7)	3 (5.1)	0.061
Anterior segments, n (%)	103 (28.8)	83 (27.8)	20 (33.9)	<0.0005
Polar map areas with reduced tracer uptake *				
Total area, (%)	3.4±4.7	5.4±5.3	0.6±1.3	0.001
Septal segments, (%)	8.3±7.3	11.5±7.2	3.6±4.4	<0.0005
Inferior segments, (%)	8.1±7.2	11.6±7.0	2.8±3.0	<0.0005
Lateral segments, (%)	2.0±3.9	3.0±4.7	0.5±1.4	0.046
Anterior segments, (%)	7.8±6.1	10.6±6.2	3.8±3.3	<0.0005